Canadian Data Report of Fisheries and Aquatic Sciences

These reports provide a medium for filing and archiving data compilations where little or no analysis is included. Such compilations commonly will have been prepared in support of other journal publications or reports. The subject matter of Data Reports reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries management, technology and development, ocean sciences, and aquatic environments relevant to Canada.

Numbers 1-25 in this series were issued as Fisheries and Marine Service Data Records. Numbers 26-160 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Data Reports. The current series name was changed with report number 161.

The correct citation appears above the abstract of each report.

Rapport statistique canadien des sciences halieutiques et aquatiques

Ces rapports servent de base à la compilation des données de classement et d'archives pour lesquelles il y a peu ou point d'analyse. Cette compilation aura d'ordinaire été préparée pour appuyer d'autres publications ou rapports. Les sujets des Rapports statistiques reflètent la vaste gamme des intérêts et politiques du Ministère des Pêches et des Océans, notamment gestion des pêches, techniques et développement, sciences océaniques et environnements aquatiques, au Canada.

Les numéros 1 à 25 de cette série ont été publiés à titre de Records statistiques, Service des pêches et de la mer. Les numéros 26-160 ont été publiés à titre de Rapports statistiques du Service des pêches et de la mer, Ministère des Pêches et de l'Environnement. Le nom de la série a été modifié à partir du numéro 161.

Le titre exact paraît au haut du résumé de chaque rapport.

Fisheries and Oceans
Canadian Data Report of
Fisheries and Aquatic Sciences
No. 395

April 1983

CATALOGUE OF SALMON STREAMS AND SPAWNING ESCAPEMENTS OF STATISTICAL AREA 4

by

M.J.Hancock, A.J.Leaney-East and D.E.Marshall

Fisheries and Oceans
Enhancement Services Branch
1090 West Pender St.
Vancouver, B.C. V6E 2P1

Minister of Supply and Services Canada 1983

Cat. No. Fs 97-13/395 ISSN 0706-6465

Correct citation for this publication:

Hancock,M.J.,A.J.Leaney-East and D.E.Marshall.1983.Catalogue of salmon streams and spawning escapements of Statistical Area 4

ABSTRACT

Hancock,M.J.,A.J.Leaney-East,and D.E.Marshall,1983. Catalogue of Salmon Spawning Streams and Spawning Escapements of Statistical Area 4 including Coastal Streams and Lower Skeena River

Catalogue containing each stream's location,spawning distribution, barriers and points of difficult ascent, escapement records and other general data pertaining to the stream. The catalogue also includes a topographical map of the stream's location and in some cases a sketch which further describes the surrounding area.

Keywords: British Columbia,Statistical Area 4,Lower Skeena River, salmon streams,spawning escapements.

RESUME

Hancock, M.J. and A.J.Leaney-East.1983. Catalogue of salmon streams and spawning escapements of Statistical Area 4 including Coastal Streams and Lower Skeena River. Can.Data Rep.Aquat. Sci.395: xxi plus 422pp.

Le présent répertoire porte sur l'emplacement de chaque cours d'eau, la répartition de fraie, les obstacles et les points de remonte difficile, les données sur les saumons de remonte et d'autres informations relatives au cours d'eau. On y trouve une carte topographique de l'emplacement du cours d'eau et, dans quelques cas, un croquis décrivant la zone environnante.

Mots-clés: Colombie-Britannique, zone statistique 4, cours inférieur de la Skeena, cours d'eau à saumons, saumons de remonte

CONTENTS

Abstract/Résumé iii
Table of Contents .... v-vi
Statistical Areas viii
Management Divisions ix
Map References x
Conservation Districts xi
Federal Fisheries Districts xii
Standards Used on Stream Data Page .... xiii-xiv
Federal Fisheries Districts Addresses xvi
Escapement Record for Statistical Area 4 xvii
Index to Map of Salmon Spawning Streams of Statistical Area 4 .... xviii-xxi

STREAM DATA

ALASTAIR LAKE 1
ALWYN CREEK 5
ANDALAS CREEK (Andulus Creek) 9
(Anderside Creek) 13
BIG FALLS CREEK 17
BURDICK CREEK (Burdock, Stoney Creek).. 21
CEDAR RIVER (Beaver River) 25
CHICAGO CREEK (Seeley Creek) 29
CHIMDEMASH CREEK 33
CLEAR CREEK 37
CLEARWATER CREEK 41
(Club Creek - Lower) 45
(Club Creek - Upper) 49
COLDWATER CREEK 53
COMEAU CREEK (Carnaby Creek) 57
CULLON CREEK (Grouse Creek) 61
DATE CREEK (Dale Creek) 65
DEEP CREEK 69
DENISE CREEK 73
(Diana Creek) 79
DOUGLAS CREEK (Dry Creek) 85
ECSTALL RIVER 89
ERLANDSEN CREEK 95
EXCHAMSIKS RIVER 99
EXTEW RIVER 103

(Extew Slough) 107
(Falls Creek) 111
(Fiddler Creek) 115
GITNADOIX RIVER 119
GLACIER CREEK 123
(Glen Vowell Creek) 127
GOAT CREEK 131
HAYWARD CREEK (Beaver Creek) 135
HAZELTON CREEK (Cedar Creek) 141
(Herman Creek) 145
(Heavenor Creek or McQueen Creek) 149
(Hot Springs Slough) 153
HUMPBACK CREEK (Humpback Bay Creek) 157
JOHNSTON CREEK 163
JOHNSTON LAKE 167
KADEEN CREEK 171
KASIKS CREEK 175
KHYEX RIVER 179
KITSEGUECHLA RIVER 183
KITSUMKALUM RIVER - LOWER 187
KITSUMKALUM RIVER - UPPER (Beaver River) 191
KITSUMKALUM LAKE 195
KITWANGA RIVER (Kitwancool) 199
KLEANZA (Gold Creek) 205
KLOIYA RIVER (Cloyah Creek) 211
KWINITSA CREEK 217
(Lahou Creek, Pearl Harbour Creek) 221
LAKELSE RIVER 227
LEAN-TO CREEK 233
(Lockerby Creek) 237
LOWRIE CREEK (Lowery Creek) 243
McCULLY CREEK 247
McNEILL RIVER (Green River) 253
McNICHOL CREEK 257
MADELINE CREEK (Gibraltar Creek) 261
MAGAR CREEK (Dog Tag Creek) 267
MOORE COVE CREEK 271
MUDDY CREEK 277
MURDER CREEK (Beirness Creek, Beirnes Creek) 281
OONA RIVER 287

(Price Creek) 293
SCHULBUCKHAND CREEK 299
SCOTIA RIVER 305
SHAMES RIVER 309
SHAMES SLOUGH 315
SHANNON CREEK (Nicholson Creek) 317
SHAWATLAN CREEK 321
SHEGUNIA CREEK 327
SILVER CREEK 333
SINGLEHURST CREEK (Swede Creek) 337
SKEENA RIVER - MAINSTEM 341
SOCKEYE CREEK 345
SOUTHEND CREEK 351
SPARKLING CREEK (Clearwater Creek) 357
SPILLER RIVER 361
STAR CREEK 367
STEPHENS CREEK 371
STUMAUN CREEK (Simpson Creek) 375
(Tsamspanaknok Bay Creek or Sam Bay)............................... 379
USELESS CREEK (Big Useless Creek) 383
USELESS CREEK (Little Useless Creek)............................... 389
WHITE CREEK 393
WILLIAMS CREEK 397
WILSON CREEK 403
ZYMAGOTITZ RIVER 407
ZYMOETZ RIVER 411

APPENDIX 1 415-421
METRIC EQUIVALENTS 422

STATISTICAL AREAS
FISHERIES & OCEANS
PACIFIC REGION


1
2A-East
2A-West
2B-East
2B-West
QUEEN
CHARLOTTE
ISLANDS
3
PRINCE RUPERT
4
5
BABINE LAKE
6
7
8
OWIKENO LAKE
RIVERS INLET
9
SMITH INLET
10
11
12
JOHNSTONE STRAIT
13
14
15
16
VANCOUVER
ISLAND
STRAIT OF GEORGIA
VANCOUVER
28
29
17
18
19
20
VICTORIA
21
22
23
24
25
26
27
JUAN DE FUCA STRAIT



MAP REFERENCES


Roads
hard surface, all weather
more than 2 lanes
hard surface, all weather
2 lanes
Route No.
18
less than 2
loose surface, all weather
2 lanes wide or more
" less than 2 lanes
all weather
dry weather
Private Road, Trail
Private Road
Trail
Railways
normal gauge, multiple track
Station
normal gauge, single track
Stop
Siding
abandoned, or under construction
narrow gauge, single track
Bridge, underpass or overpass
Tunnel


Boundary, International
" Province
" County or District
" Township or Parish
" City or Town
" Reservation, Indian, Military, etc
Power Transmission Line
Telephone or Telegraph, trunk route
Horizontal Control Point
Boundary Marker
Bench Mark
BM
2975
Spot Elevation, (in feet)
5752
Mine or Pit


Road, Hard Surface, All Weather
Route No.
99
2 Lanes
" Loose Surface, All Weather
2 Lanes
" Loose Surface, Less than 2 lanes
All Weather
Dry Weather
" Private (Logging, Mining etc
" Four Wheel Drive
Trail
Railway
Main Telephone Line
Main Electric Power Line
Horizontal Control Station
Contours (Interval 500 feet)
5000
Elevation in feet above mean sea-level
2584 (6312)
Intermittent Stream
Swamp or Marsh
Dam
Dam
Spring
S
Navigation Light
Mine
Glacier
Customs Office
Cust


House, Building
School
S
Church
" with conspicuous Tower or Spire
Post Office
P
Tower, Radio Mast, Lookout, etc
Cemetery
Cem
Quarry
Sand or Gravel Pit
Cliff
Cutting
Embankment
Saw Mill
SM


Lighthouse
Wharf or Pier
Foreshore Flats
Sand
Mud
Swamp or Marsh
Lake or Pond, intermittent
Glacier or Snowfield
Stream, intermittent
Irrigation Canals, Ditches
Inundated Land, seasonal
Contours, elevation
500
400
" depression
500
400
" approximate
500
400
Forest, unclassified


Surveyed timber license number
TL 2841
Lot number
L 124 or S 66
Building
School
Non-perennial stream
Marsh or Swamp
Glacier
Foreshore flats
Sand
Contours, elevation
500
Contours, depression
Forest


City or large town
Town
Village or settlement
Post office
P
School
Church
Boundary monument
Astronomical position
Horizontal control point
Streams
intermittent or dry
indefinite
Irrigation canal or ditch
Rapids, falls
R
F
Aerodrome
Landing ground
Intermittent lake
Marsh or swamp
Sand, gravel or mud
Wooded areas
Seaplane base
Seaplane anchorage


Streams
Highways
Roads
Trails
Houses
Railroad
Falls
Rapids
Rip-Rap
Bridges
Dam
Log Jams
Log
Power Line
Coho
CO
Chum
CM
Pink
P
Chinook
CK
Sockeye
S



CONSERVATION DISTRICTS
FISHERIES & OCEANS
PACIFIC REGION

YUKON
TERRITORY
BRITISH COLUMBIA
C.D. 10
C.D. 8
C.D. 9
QUEEN
CHARLOTTE
ISLAND
C.D. 7
C.D. 1
C.D. 5
VANCOUVER
ISLAND
C.D. 2
C.D. 3
C.D. 4
C.D. 6



FEDERAL FISHERIES DISTRICTS

FISHERIES & OCEANS PACIFIC REGION

LEGEND
DISTRICT BOUNDARY
SUBDISTRICT BOUNDARY
★ DISTRICT OFFICE
☆ SUBDISTRICT OFFICE
YUKON
NORTHERN B.C.
NASS RIVER
TERRACE
SMITHERS
LOWER SKEENA
MASSET
GRENVILLE - PRINCIPE
KITIMAT
EAST COAST
WEST COAST
PRINCE GEORGE
BELLA COOLA
BELLA BELLA
CARIBOO
CLEARWATER
RIVERS INLET
SMITH INLET
SEYMOUR INLET
ALERT BAY
CAMPBELL RIVER
LILLOOET
SALMON ARM
QUATSINO SOUND
KYUQUOT
TAHSIS
TOFINO
PORT ALBERNI
POWELL RIVER
HOWE SOUND
MADEIRA PARK
MISSION
CHILLIWACK
NEW WESTMINSTER
VANCOUVER
STEVESTON
NANAIMO
VICTORIA
HOWE SOUND
VANCOUVER
MISSION
NEW WESTMINSTER
STEVESTON
CHILLIWACK

STANDARDS USED ON STREAM DATA PAGE

Name of Stream: Name as given in Gazetteer of Canada, British Columbia 1966 edition; local or non-gazetted names are added in lower case type.

Conservation District: As defined by the Conservation and Protection Service (April 1965).

Statistical Area: As defined by Department of the Environment, Fisheries Operations Statistical Map (January 1974).

Location and Position: Defined by quadrant indexing. Each geographical quadrilateral of the earth's surface of 1 degree in extent in latitude and longitude is divided into the SE, SW, NE and NW quarters. The south-east corner of each quadrilateral gives the initial point for the figure of reference (Gazetteer of Canada).

EXAMPLE "X"
53° 122° NW



Length: The portion of the stream accessible to spawning salmon.

Width: Average width, estimated to the nearest metre for the described length.

Drainage: Area in square kilometres of the entire drainage basin feeding the stream.

Composition:

Bedrock	bedrock
Boulder	>256 mm
Coarse	50.9 - 256 mm
Fine	3.37 - 50.8 mm
Sand & Silt	<3.37 mm
Unclassified	where bottom cannot be observed, e.g. log jams, pools, water colour, etc.

Gradient: Expressed as a percentage

Wetted Area: Number of square metres of stream bed under water at average flows within the described length.

Spawning Area: Estimated square metres of stream bed suitable for salmon spawning within the described length.

Discharge: Mean annual discharge. Maximum and minimum values are either daily means or instantaneous discharges. The latter are identified by (Inst.). Discharge data is taken from "Historical Stream Flow Summary", British Columbia, Water Survey of Canada.

Temperature: As described. (°C)

Barriers and Points of Difficult Ascent: Complete and partial barriers to salmon and their distance from the stream mouth. Species likely to be affected may be listed. Both natural and man-made obstructions are defined.

Spawning Distribution: Portion of the stream utilized by each species. Distribution is indicated by brief comments opposite the species.

Potential of Inaccessible Portion of Stream: Indicates whether or not the inaccessible portion of the stream could be utilized by spawning salmon.

General Remarks: Emphasizes features of stream and spawning populations. Also includes industrial activity, routes of accessibility, etc. The comments with dates are taken from "Annual Reports of Salmon Streams & Spawning Grounds" (B.C. 16's). In some cases, references to additional information not included in the General Remarks may be given.

Escapement Records: The escapement represents the mid point of the coded range of escapement for each species. For example: the letter "H" representing 5000-10000 fish would be entered as 7500. Where absolute numbers are provided by Fisheries Personnel, these numbers are entered. N/O means the stream was inspected but no fish were observed; UNK means there was evidence of fish present but no estimates were made; NO RECORDS means no escapement records for the applicable years could be found in the escapement files.

Timing: Dates which salmon arrive in the stream, begin to spawn, reach peak spawning period and finish spawning.

E = early (1st to 10th of the month)
M = mid (11th to 20th of the month)
L = late (21st to end of the month)

NB: Distance references are from the mouth of the stream unless otherwise stated.

BRITISH COLUMBIA
FEDERAL FISHERIES DISTRICTS

DISTRICT AND SUBDISTRICT OFFICES	ADDRESSES	TELEPHONE NUMBERS
DISTRICT #1 KAMLOOPS	202 Federal Bldg., 317 Seymour St., Kamloops, B.C., V2C 2E9	374-4322
Salmon Arm	Box 1160, Salmon Arm, B.C., V0E 2T0	832-8037
Prince George	Box 267, Postal Stn. A, Prince George, B.C., V2L 4S1	564-7030
Clearwater	P.O. Box 610, P.O. Building, Clearwater, B.C., V0E 1N0	674-9633
Lillooet	Box 315, Lillooet, B.C., V0K 1V0	256-4525
Cariboo	Box 4340, Quesnel, B.C., V2J 3J3	992-2434
DISTRICT #2 NEW WESTMINSTER	309 - 549 Columbia St., New Westminster, B.C., V3L 1B3	545-7181
Surrey	309 - 549 Columbia St., New Westminster, B.C., V3L 1B3	545-7181
Vancouver Waterfront	309 - 549 Columbia St., New Westminster, B.C., V3L 1B3	545-7181
Mission	Box 3308, Mission, B.C., V2V 4J5	826-3664
Chilliwack	Ste. 5, 9 Mary Street, Chilliwack, B.C., V2P 4G9	792-6011
Steveston	1255 No. 1 Road, Richmond, B.C., V7E 1T7	274-7217
Coquitlam	309 - 549 Columbia St., New Westminster, B.C., V3L 1B3	545-7181
Squamish	Box 85, Squamish, B.C., V0N 3G0	892-3230
DISTRICT #3 NANAIMO	204 - 60 Front St., Nanaimo, B.C., V9R 5H7	876-274
Parksville	Box 1120, Parksville, B.C., V0R 2S0	248-6710
Comox	Box 1328, Comox, B.C., V9N 3Z9	339-2031
Duncan	Box 241, Duncan, B.C., V9L 3X3	746-6221
Powell River	4770 Joyce Ave., Powell River, B.C., V8A 3B6	485-9621
Pender Harbour	Box 10, Madeira Park, B.C., V0N 2H0	883-2313
DISTRICT #4 PORT ALBERNI	P.O. Box 280, Federal Bldg., Port Alberni, B.C., V9Y 7M7	724-0195
Tofino	Box 48, Tofino, B.C., V0R 2Z0	725-3468
Tahsis	Box 549, Tahsis, B.C., V0P 1X0	934-6606
Port Hardy	Box 10, Port Hardy, B.C., V0N 1P0	949-6422
DISTRICT #5 CAMPBELL RIVER	215 - 950 Alder St., Campbell River, B.C., V9W 2P8	287-2102
Port Hardy	Box 10, Port Hardy, B.C., V0N 2P0	949-6422
Alert Bay	Box 263, Alert Bay, B.C., V0N 1A0	974-5216
DISTRICT #6 VICTORIA	116 - 816 Government St., Victoria, B.C., V8W 1W9	566-3252
Saanich	116 - 816 Government St., Victoria, B.C., V8W 1W9	566-3252
Sooke	Box 831, Sooke, B.C., V0Z 1N0	642-5322
DISTRICT #7 KITIMAT	312 - 450 Federal Bldg., City Centre, Kitimat, B.C., V8C 1T6	632-6158
Butedale	312 - 450 Federal Bldg., City Centre, Kitimat, B.C., V8C 1T6	632-6158
Bella Bella	Box 38, Bella Bella, B.C., V0T 1B0	957-2312
Bella Coola	Box 130, Bella Coola, B.C., V0T 1C0	799-5345
Rivers Inlet	Dawsons Landing P.O., Rivers Inlet, B.C., V0N 1M0	Radio Ph.
DISTRICT #8 PRINCE RUPERT	109 - 417 Second Ave. West, Prince Rupert, B.C., V8J 1G8	624-9137
Lower Nass	109 - 417 Second Ave. West, Prince Rupert, B.C., V8J 1G8	624-9137
Upper Nass	P.O. Box 29, Nass Camp P.O., B.C., V0N 3J0	633-2408
Skeena	109 - 417 Second Ave. West, Prince Rupert, B.C., V8J 1G8	624-9137
Grenville-Principe	109 - 417 Second Ave. West, Prince Rupert, B.C., V8J 1G8	624-9137
Hazelton	Box 327, Hazelton, B.C., V0J 1Y0	842-6327
Terrace	4721-B Lazelle Ave., Terrace, B.C., V8G 1R6	635-2206
Smithers	Box 578, Smithers, B.C., V0J 2N0	847-2312
DISTRICT #9 QUEEN CHARLOTTE	Box 99, Q.C. City, Queen Charlotte Islands, B.C., V0T 1S0	559-4413
Sandspit	Box 222, Sandspit, B.C., V0T 1T0	637-5340
Masset	Box 99, Masset, B.C., V0T 1M0	626-3316
DISTRICT #10 WHITEHORSE	122 Industrial Road, Whitehorse, Yukon Territories, Y1A 2T9	403-667-2235
Haines Junction	P.O. Box 5341, Haines Junction, Y.T., Y0B 1L0	403-667-2235

ESCAPEMENT RECORD FOR STATISTICAL AREA 4 LOWER SKEENA RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947	31,254	4,475	14,125	10,850	86,400	25
1948	55,425	1,575	19,850	18,875	248,000	-
1949	43,960	2,341	41,795	12,130	470,960	-
1950	44,600	3,075	20,042	24,450	476,875	-
1951	40,300	5,430	37,815	16,500	396,575	-
1952	74,290	3,000	28,679	31,175	700,000	-
1953	44,550	2,125	28,900	8,165	263,600	-
1954	49,475	3,750	38,925	22,997	314,095	-
1955	31,625	4,825	42,550	2,075	294,050	-
1956	63,430	4,850	76,687	3,820	189,384	26
1957	68,675	10,325	44,875	8,525	521,450	-
1958	79,207	6,050	101,100	1,945	363,090	75
1959	46,875	9,375	875	1,225	537,525	-
1960	25,882	6,852	40,500	600	180,185	-
1961	36,352	7,475	39,568	37,562	727,450	59
1962	29,537	6,125	18,750	8,200	874,575	133
1963	35,600	6,900	39,675	1,325	817,075	75
1964	60,129	6,050	59,375	3,050	1,343,203	-
1965	56,880	12,131	108,750	8,653	1,224,859	-
1966	37,306	5,858	102,138	18,580	594,313	-
1967	43,336	5,100	44,925	13,775	472,077	243
1968	43,875	9,950	96,500	10,300	1,613,400	-
1969	24,625	19,000	57,600	10,225	299,425	-
1970	25,900	12,725	61,675	10,475	784,300	-
1971	25,400	11,475	57,745	24,000	608,781	-
1972	30,625	7,430	41,200	33,620	35,000	-
1973	27,450	13,770	24,120	23,895	722,800	-
1974	16,700	12,000	23,920	12,400	344,700	-
1975	13,500	11,075	15,420	9,425	1,216,105	-
1976	12,950	13,330	25,820	10,245	574,700	-
1977	20,175	22,575	20,120	10,125	569,925	200
1978	21,400	15,645	25,350	7,175	523,675	-
1979	16,075	2,300	11,550	6,175	354,195	-
1980	23,632	17,147	15,343	3,781	155,152	-
1981	15,595	3,725	7,785	8,910	41,495	-
1982						
1983						
1984						
1985						
TIME						
ARR.						
START						
PEAK						
END						

REMARKS


Salmon Spawning Streams
of Lower Skeena River
0
24
kilometres
N
Terrace
SKEENA RIVER
Prince Rupert
Porcher
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
17
18
19
21
22
23
25
26
27
28
31
33
34
35
36
37
38
39
41
42
43
44
45
47
51
57

INDEX TO MAP OF SPAWNING STREAMS OF STATISTICAL AREA 4
LOWER SKEENA RIVER

ALASTAIR LAKE	(38)
ALWYN CREEK	(49)
ANDALAS CREEK	(44)
(Anderside Creek)	(33)
BIG FALLS CREEK	(24)
CLEARWATER CREEK	(45)
COLDWATER CREEK	(42)
DEEP CREEK	(56)
DENISE CREEK	(9)
(Diana Creek)	(6)
ECSTALL RIVER	(23)
ERLANDSEN CREEK	(57)
EXCHAMSIKS RIVER	(30)
EXTEW RIVER	(31)
(Extew Slough)	(34)
GITNADOIX RIVER	(35)
GLACIER CREEK	(64)
GOAT CREEK	(65)
HAYWARD CREEK (Beaver Creek)	(25)
(Herman Creek)	(48)
(Hot Springs Slough)	(47)
HUMPBACK CREEK (Hampback Bay Creek)	(3)
JOHNSTON CREEK	(21)
JOHNSTON LAKE	(22)
KADEEN CREEK	(36)
KASIKS CREEK	(29)
KHYEX RIVER	(16)
KITSUMKALUM RIVER - LOWER	(62)
KLUIYA RIVER (Cloyah Creek)	(7)
KWINITSA CREEK	(28)
(Lahou Creek, Pearl Harbour Creek)	(13)
LAKELSE RIVER	(92)
(Lockenby Creek)	(26)
McNEILL RIVER (Green River)	(8)
McNICHOL CREEK	(10)
MADELINE CREEK (Gibraltar Creek)	(18)
MAGAR CREEK (Dog Tag Creek)	(39)
MOORE COVE CREEK	(17)
MUDDY CREEK	(19)
OONA RIVER	(4)
SCHULBUCKHAND CREEK	(46)
SCOTIA RIVER	(27)
SHAMES RIVER	(40)
(Shames Slough)	(41)
SHAWATLAN RIVER	(11)
SILVER CREEK	(12)
SKEENA RIVER - MAINSTEM	(32)
SOCKEYE CREEK	(50)
SOUTHEND CREEK	(37)
SPARKLING CREEK (Clearwater Creek)	(20)
SPILLER RIVER	(5)
STAR CREK	(63)
STUMAUN CREEK (Simpson Creek)	(14)
(Tsamspanaknok Bay Creek, Sam Bay)	
USELESS CREEK (Big Useless Creek)	
USELESS CREEK (Little Useless Creek)	
WHITE CREEK	(43)
WILLIAMS CREEK	(51)
ZYMAGOTITZ RIVER	(63)


Salmon Spawning Streams
of Lower Skeena River
N
0
24
kilometres
91
90
89
88
Kispiox
87
River
86
85
83
84
82
80
81
79
Hazelton
78
76
75
77
72
73
74
SKEENA RIVER
69
70
71
68
66
67
Kitsumkalum
Lake
61
59
52
55
53
54

INDEX TO MAP OF SPAWNING STREAMS OF STATISTICAL AREA 4

LOWER SKEENA RIVER

BURDICK CREEK (Burdock, Stoney Creek)	76
CEDAR RIVER (Beaver River)	69
CHICAGO CREEK (Seeley Creek)	78
CHIMDEMASH CREEK	55
CLEAR CREEK	70
(Club Creek - Lower)	89
(Club Creek - Upper)	90
COMEAU CREEK (Carnaby Creek)	77
CULLON CREEK (Grouse Creek)	86
DATE CREEK (Dale Creek)	82
DOUGLAS CREEK (Dry Creek)	71
(Falls Creek)	91
(Fiddler Creek)	66
(Glen Vowell Creek)	80
HAZELTON CREEK (Cedar Creek)	79
(Heavenor Creek, McQueen Creek)	84
KITSEGUECLA RIVER	74
KITSUMKALUM RIVER -- UPPER (Beaver River)	68
KITSUMKALUM LAKE	67
KITWANGA RIVER (Kitwancool)	75
KLEANZA CREEK (Gold Creek)	53
LEAN-TO CREEK	60
LOWRIE CREEK (Lowery Creek)	59
McCULLY CREEK	83
MURDER CREEK (Beirness Creek, Beirnes Creek)	85
(Price Creek)	73
SHANNON CREEK (Nicholson Creek)	61
SHEGUNIA CREEK	81
SINGLEHURST CREEK (Swede Creek)	52
STEPHENS CREEK	88
WILSON CREEK	72
ZYMOETZ RIVER	54

STREAM DATA

STATISTICAL AREA 4 (Lower Skeena River)


N
Alastair Lake
0
2
kilometres
MOUNTAINS
FOREST
ALASTAIR LAKE
VL
VK
ALASTAIR
IR 80
L 7184
ALASTAIR
IR 82
L 7183
ALASTAIR
Alastair Peak
84
85
86
87
88
89
90
91
92
00
99
98
97
96
95
94
93
92
91
90
89
4300
2500
4000
1400
5200
2700
4700
4000
5800
2100
1900
5600
4800
1800
5600
4200
700
2000
2800

NAME OF STREAM ALASTAIR LAKE

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Expansion of Gitnadoix River, Rge.5, Coast Dist.

POSITION 54 129 SE

LENGTH km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT 15.2m falls 1.6km from mouth

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- scattered along lake shore and at creek mouths to a depth of
CHINOOK	4-6m
COHO	- tributary streams
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS

1960 - Good seeding of sockeye in creeks above lake.

1964 - Usual silty output off Southend Creek sandy lake edge.Fish ripen at lake edge. Bad weather made late season observations impossible.

Predators: seals, bears, eagles and other birds.

Subject to seasonal fluctuations in water levels.

Lake fed by two small streams with adequate water volume and cool temperatures.

Stable lake conditions.


Sketch of Alastair Lake,
1971.
N
0
1.50
kilometres
Alastair Lake
× - Sockeye Spawning Areas
□ - Coho Spawning Areas
Gitnadoix
River
Kadeen Creek

ESCAPEMENT RECORD FOR ALASTAIR LAKE

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947	7500					
48						
49	2000					
50	3000					
51	3000					
52	5000					
53	3000					
54	3500		* 7000			
55	3500		* 6000			
56	7500					
57	7500					
58	7500		* 7500			
59	7500		N/I			
60	400		N/I			
61	750					
62	1500		N/I			
63	1500					
64	400		750			
65	400					
66	400					
67	5000		500			
68	7500		1500			
69	1500		1500			Obs.
70	1500		1500			
71	400		750			
72	1500		750			
73	1500		400			
74	1500		400			
75	200		400			
76	1500		750			
77	3500		200			
78	1500		400			
79						
80	4500		200			
81	500		300			
82						
83						
84						
85						

TIMING

ARRIVE	E AUG		SEPT			
START	E-M SEPT		OCT			
PEAK	M SEPT		M NOV			
END	OCT		DEC			

REMARKS * Sighted in lake,but spawning location unknown.
N/I - Not inspected
Obs - Observed


Alwyn Creek
N
0
2
kilometres
SKEENA
SKEENA RIVER
HELLSGATE
SLOUGH
Thunderbird
Alwyn
Creek
ZYMAGOTITZ
R
ZIMAGORD IR 3
Kulspai IR 6
Terrace
LAKELSE
T L 9509
BLK A
16

NAME OF STREAM ALWYN CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows W. into Skeena R., Rge.5 Coast Dist.

POSITION 54 128 SW.

LENGTH 8 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- 3-8 km upstream from mouth
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS Log jams at approx 8 km from mouth of stream

1970 Winter Works Programme cleaned out log jams in lower 3 km. Recommended work in upper section before stream is suitable for salmon propagation.

1971 Recommended removal of beaver dams.

1980 Numerous fry observed.

Heavy silting in lower 1.6 km due to logging debris -- some scouring (20%) after heavy rainfall.

Normal water levels

No predators observed except for 1980 when a heavy bear population in area.



◘ – Coho spawning grounds
Logging Road
To Terrace
Alwyn Creek
SKEENA RIVER
N
0
2.0
kilometres

Sketch of Alwyn Creek, 1971.

ESCAPEMENT RECORD FOR ALWYN CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70			25			
71			25			
72			25			
73			25			
74			25			
75			25			
76			25			
77			25			
78			75			
79			N/R			
80			25			
81			50			
82						
83						
84						
85						

TIMING						
ARRIVE						
START			OCT			
PEAK			E OCT			
END			M NOV			
			DEC			

REMARKS N/R - No record

Two Chinook observed in 1980


Andalas Creek
N
0
2
kilometres
Squirrel Pt
Lakelse Hot Springs
Campsite
Ena Lake
Ena Ck
Andalas Ck
Clearwater Creek
TREE FARM LICENCE 41
FORCEMAN RIDGE
Schulbuckhand
West Lake
Onion Lake
Clearwater Lakes
Cecil Creek
Du Bose
CANADIAN NATIONAL
2662 T L 8288
2663 T L 11145
2664 T L 11144
2665 T L 11146
2672 T L 11135
2674 T L 8276
2675 T L 8277
2676 T L 11370
2677 T L 11129
2678 T L 11128
2679 T L 11127
2680 T L 11126
2684 T L 8278
2688 T L 11130
2689 T L 11363
2690 T L 11362
2691 T L 11361
2693 T L 8279
2695 T L 11143
T L 7723
T L 7724
3983
3988
3987
5127
5129
5139
5140
5965
6794
6800
6801
6802
6803
6804
6805
6806
6807
6808
6809
6891
684
700
1000
500
25

NAME OF STREAM ANDALAS CREEK (Andulus Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows NE. into S. end of Lakelse Lake, Rge., 5 Coast Dist.

POSITION 54 128 SW

LENGTH km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

Passable beaver dams at various distances from stream mouth.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- most spawning occurs on 183 m of gravel located 0.8 km above stream mouth
CHINOOK	
COHO	- from mouth to 0.8 km
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1949 - Fish moved through system over a two month period.

1960 - Forest debris 2.4 km from mouth. Suggested removal as required.

1962 - Late coho was not inspected.

1964 - 1968 and 1970 - 1981 Removed beaver dams in low marshy region of stream as required.

1965 - Winter logging expected to proceed in vicinity.

1967 - Marshy area functions as rearing for Lakelse Lake stock.

Small stream -- muddy bottom for first 1.6 km.

Subject to seasonal fluctuations in water levels.

Predators: bears and eagles.



Sketch of Andalas Creek, 1974.

Scully Creek
Lakelse Lake
Clearwater Creek
Coho
Sockeye
Andulus Creek
N
0
.48
kilometres

ESCAPEMENT RECORD FOR ANDALAS CREEK (Andulus Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48			200			
49			200			
50			200			
51			70			
52			100			
53			100			
54			25			
55			75			
56			75			
57			75			
58			200			
59			200			
60			N/I			
61			400			
62			N/I			
63			N/I			
64			200			
65			500			
66	25		750			
67	N/O		50			
68	25		400			
69	150		300			
70	25		25			
71	25		25			
72	25		25			
73	25		25			
74	25		25			
75	25		25			
76	25		25			
77	25		75			
78	25		75			
79	25		75			
80	20		225			
81	20		100			
82						
83						
84						
85						

TIMING

ARRIVE	E AUG		E OCT			
START	M-L AUG		M OCT -E NOV			
PEAK	L AUG -E SEP		M NOV			
END	M-L SEPT		M NOV - DEC			

REMARKS N/I - Not inspected


Andeside Creek
N
0
2
kilometres
RANGES
HUDSON
BAY
FLATS
Extew Slough
Andeside Creek
CANADIAN NATIONAL
Exstew
RIVER
L 5631
L 7176
Block B
L 5626
Block C
Block A
L 540
L 3999
L 5628
TL 9503
L 5866
TL 9504
16
2000
1500
700
1800
100
3500
4000
2500
3000
88
89
90
91
93
94
95
96
97
98
99
21
24
25
26
27
28
29
30
31

NAME OF STREAM (Anderside Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows into Skeena River, near Extew Slough.

POSITION 54 129 SE.

LENGTH 2 km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT cascading rapids 2km from mouth.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	- in slough below C.N.R.Bridge.
PINK (ODD YEAR)	- " "
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1965 Minor stream but supplies slough with good flow of cold, clear mountain water,and fish spawn in slough below stream.

1966 - Swift upper reaches,steep rocky contours. Flood controls channelled.

1967 - Annual flash flood, scouring, sand and silt .

1969-70 Water levels fluctuate seasonally.

1974 - Heavy erosion, silting and scouring during Autumn floods,1975 scouring near mouth.

Predators: light, a few eagles.


Extew Station
Hudson Bay
Flats
▨ – Pink and Chum spawning grounds
Andeside Creek
N
0
2
kilometres
SKEENA RIVER

Sketch of Anderside Creek, 1971

ESCAPEMENT RECORD FOR (Anderside Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63.						
64						
65						
66				400	400	
67				200	75	
68				100	200	
69				400	400	
70				200	200	
71				750	200	
72				400	200	
73				1500	750	
74				750	200	
75				400	25	
76				200	-	
77				200	400	
78				25	25	
79				800	50	
80				-	-	
81				150	25	
82				20	200	
83						
84						
85						

TIMING						
ARRIVE				Aug	Aug	
START				Aug-L Aug	EAug-M Aug	
PEAK				LAug-M Sept	ESept-LSept	
END				Sept	Sept-Oct	

REMARKS


N
Big Falls Creek
0
10
kilometres
TREE FARM LICENCE 1
BLOCK 1
Telegraph Point
Carnation
Tyee
Raspberry Is
Port Essington
Ecstall
BALMORAL PK
SKIP MTN
Windsor Pt
Aberdeen
Khtada Lake
Kwinitsa
Salvus
MT HAYWARD
Big Falls
Herman L
Madeline L
Ecstall River
Brown Lake
Muddy
Kumealon L
Kumealon Lagoon
McMurray Pt.
Baker Inlet
Griffon Pt.
Johnston L
MT MORRISON
SPOKSHUTE MTN
Alder
R A N
F O
K I

NAME OF STREAM BIG FALLS CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows W. into Ecstall R., Rge.4,Coast Dist.

POSITION 53 129 NW

LENGTH ____ km WIDTH ____ m DRAINAGE ____ km^2

COMPOSITION: BEDROCK ____ BOULDER ____ COARSE ____ FINE ____

SILT & SAND ____ UNCLASSIFIED ____

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ____ m^2 SPAWNING AREA ____ m^2

DISCHARGE (m^3/s) ____

TEMPERATURE (°C) ____

BARRIERS OR POINTS OF DIFFICULT ASCENT Power dam 365m from stream mouth. Passable rock falls 91m from mouth.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- below power house
COHO	
CHUM	
PINK (ODD YEAR)	- on gravel bars below falls 274m from stream mouth.
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ____

GENERAL REMARKS 1953 Some scouring of streambed when dam sluice gates open due to turbulence. Tidal influence and power plant overflow prevent ice formation and reduces desiccation threat.

1954-1956 average weight of pink salmon ranged between 1.86 kg and 2.54kg

About 2100 square meters of spawning area. Remainder of stream silty and tidally affected.

Small spawning population. Good water quality. Good gravel deposit.

Flows controlled by power dam following 1953. Subject to seasonal fluctuations prior to 1953.

1981 D.B.Lister and Associates prepared a comprehensive study and made spawning observations for B.C.Hydro.

1976 Extremely limited spawning area below dam;possibility for enhancement is almost nil. Predators: trout and hair seals.



Sketch of Big Falls Creek.

Wharf
Float
Powerhouse
End of Tidal Influence
Power Dam
Ecstall River
Rockfall
Gravel Bars
Pink Salmon Spawning Area
N
0
.32
kilometres

ESCAPEMENT RECORD FOR BIG FALLS CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48					750	
49					400	
50					1000	
51					200	
52					300	
53					1000	
54					400	
55		75			500	
56					1500	
57					50	
58		50			75	
59		25			100	
60		50			400	
61		25			25	
62		25			50	
63					25	
64		25			750	
65		25			25	
66		25		200	400	
67		N/O		N/O	N/O	
68		50		25	200	
69		25			400	
70		N/O		200	N/O	
71		N/O		N/O	N/O	
72		50		N/O	N/O	
73		20			N/O	
74		N/O		N/O	50	
75		N/O			N/O	
76		N/O			N/O	
77		N/O			N/O	
78		N/O			N/O	
79		25			N/O	
80		27				
81						
82						
83						
84						
85						

TIMING

ARRIVE	M AUG					
START	L AUG				AUG	
PEAK	E SEPT				SEPT	
END	L SEPT-OCT				OCT	

REMARKS N/O - None observed


SKEENA PROVINCIAL FOREST
RANGE
SKEENA RIVER
Keynton Lake
Keynton Creek
KISPIOX
N
0
2
kilometres
22
21
20
19
18
16
15
L 1620
L 1619
L 1616
L 1615
L 2151A
L 2151
L 2150
L 2147
L 2148
L 2152
L 2153
L 2926
L 2927
L 2924
L 2925
L 2928
L 2149
L 818
L 819
L 820
L 1282
L 2382
BKA
BKB
BKD
1500
1000
2000
2500
3500
Highway

Burdick Creek

For more detail see appendix p. 416

NAME OF STREAM BURDICK CREEK (Burdock,Stoney Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S.E. into Skeena River, N. of Carnaby, Cassiar Dist.

POSITION 55 127 SW

LENGTH ______ km WIDTH ______ m DRAINAGE ______ km²

COMPOSITION: BEDROCK ______ BOULDER ______ COARSE ______ FINE ______

SILT & SAND ______ UNCLASSIFIED ______

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ m²

SPAWNING AREA ______ m²

DISCHARGE (m³/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT ______

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	- from mouth to 3.2km upstream
PINK (EVEN YEAR)	- " " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ______

GENERAL REMARKS

1952,1954-58,1961 - Not observed during coho spawning season.

1960 - Bouldery substrate near bridge. Low water 1960-61 prevented fish from entering creek.

1962 - Heavy scouring destroying habitat. Rock falls 3.6m located 0.8km from mouth.

Steep terrain - 10% grade.

Subject to occasional low flows.

Predators: bears.

ESCAPEMENT RECORD FOR BURDICK CREEK (Burdock, Stoney Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			25		200	
48					200	
49			25		400	
50			75		400	
51			75		1000	
52			N/I		200	
53			–		300	
54			N/I		750	
55			N/O		750	
56			N/I		200	
57			N/I		750	
58			N/I		75	
59			N/I		N/I	
60			N/I		N/O	
61			N/I		N/O	
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81						
82						
83						
84						
85						
TIMING						
ARRIVE						
START			E SEPT			
PEAK			M SEPT			
END			L SEPT			

REMARKS 1958 Coho reported on this stream by hunters
1962 Report: No fish seen; too rocky and steep. Scoured out.

N/I - Not inspected
N/O - None observed



Cedar River

For more detail see appendix p. 417

N
0
10
kilometres
TREE FARM
LICENCE 1
BLOCK 4
KITIMAT
RANGES
Aiyansh
New Aiyansh
Canyon City
Hoodoo
MT HOADLEY
MT PRIESTLEY
MT HOEFT
MT PHILLIPPS
VETTER PK
ALDER PK
Lava
Lake
Shumal
Maughan
Chambers
Kwinamuck
River
Cr
Alder
Sand
COAST LAND DISTRICT RANGE 5

NAME OF STREAM CEDAR RIVER (Beaver River)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S. into Kitsumkalum River, N.W. of N. end of Kitsumkalum Lake, Rge. 5, Coast Dist.

POSITION 54 128 NW

LENGTH 16 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m²

SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Numerous passable log jams. Swift, bouldery canyons 26-29km upstream.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1965-66 Stream cleared. Opened up spawning area for approx 16km. Good gravel and spawning potential.

1962 -- Logging reducing timber canopy within Columbia Cellulose Management area. Protective clauses required in cutting permits to assure 40.2m of timber edge. Sub contractors operate under Columbia Cellulose direction.

1963 - Logging off watershed C.C.Co tree farm area.

1965 - Clear to canyons. Checked 32km by helicopter compliments of Columbia Cellulose Co.

continued.....

continued.

GENERAL REMARKS   CEDAR RIVER

Silting, erosion and scouring mentioned in 1967-77 reports. Frequent changes in stream course in lower section. Watershed has been extensively logged. Stream is prone to flash floods causing extensive erosion in upper section and forming new channels in lower section.

1973 - Log jams, blow downs and debris in lower 11-12km. Numerous log jams but not impassable for fish. Recommend removal as necessary.


Clarence Creek
Sterling Creek
Belway Creek
Cedar River
Beaver River
◘ - Coho Spawning Grounds
Δ - Chinook Spawning Grounds
x - Sockeye Spawning Gounds
N
0
2.56
kilometres

Sketch of Cedar River, 1971.

ESCAPEMENT RECORD FOR CEDAR RIVER 28

(Beaver River)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947	400	N/O	400			
48						
49	500	Obs	Obs			
50	Obs	Obs				
51						
52	*N/I					
53	*N/I					
54	*N/I					
55	N/O	750	N/O			
56	750	750	N/O			
57	N/O	1500	N/O			
58	750	1500	Obs			
59	N/I	N/I	N/I			
60	400	200	25			
61	400	25	75			
62	750	25	25			
63	1500					
64	3000	100	2000			
65	1500	200	1500	25		
66	1500	25	200			
67	1000	50	200			18
68	1500	25	1500			N/O
69	1500	200	200			X
70	1500	400	200			N/O
71	1500	400	1500			N/O
72	1500	400	1500			X
73	1500	750	1500			
74	750	750	1500			
75	750	750	1500			
76	400	750	1500			
77	400	4000	750			
78	500	750	1500			
79			750			
80	200	600	750			
81	100	600	700			
82						
83						
84						
85						

TIMING

ARRIVE	AUG	JULY	OCT			
START	M Aug-Sept	JUL-L AUG	OCT- E NOV			
PEAK	L Aug-L Sep	M AUG-L SEP	OCT- L NOV			M SEPT
END	L Sept-Nov	M - L SEP	OCT - DEC			L SEPT
						DEC

REMARKS * 1952/3/4 Settlers reported average to good sockeye runs

X 1968 and 1971 Anglers reported normal Steelhead run

N/I - Not inspected

N/O - None observed

Obs - Observed


Chicago Creek
N
0
2
kilometres
SKEENA
ANGES
NATIONAL
Seeley Lake
Provincial Park
Seeley
Lake
Sawmill
Burner
Theatre
Chicago
BM 937
Sawmill
Glacier
Silvertip
Glacier
Abandoned
21
20
19
18
17
16
15
14
13
12
11
88
89
90
1000
1500
2500
3000
3500
4500
5000
5500
6000
L 699A
L 699
L 778
L 892
L 890
L 1576
L 1374
L 217
L 216
L 698
L 776
L 318
L 1375
L 781
L 2126
L 1376
L 2523
L 1380
L 1379
L 1378
L 1382
L 1381
L 2384
L 4272
L 6250
L 6257

NAME OF STREAM CHICAGO CREEK (Seeley Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows W. into Skeena River N. of Seeley Lake, Cassiar Dist.

POSITION 55 127 SW

LENGTH ______ km WIDTH ______ m DRAINAGE ______ km²

COMPOSITION: BEDROCK ______ BOULDER ______ COARSE ______ FINE ______

SILT & SAND ______ UNCLASSIFIED ______

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ m² SPAWNING AREA ______ m²

DISCHARGE (m³/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT C.N.R.culvert 3.2 km from mouth is difficult to negotiate during low flow.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- to C.N.R.culvert and beyond in years of high flow.
CHUM	
PINK (ODD YEAR)	- near mouth
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ______

GENERAL REMARKS 1947 Coho had difficulty entering stream due to low water levels in Skeena.

1958 - Low water prevented coho from entering stream.

1962 - Popular campsite and trout fishing resort.

1965 - Meanders through swamps near mouth. Suggest stop logs be installed to raise water level in culvert and ease the 1m drop between culvert and water surface.

1966 - Numerous side channel rearing grounds.

1970,76,77,78 Scouring during high water and C.N.R.agreed to construct fishway at lower end of culvert to enable fish to pass upstream.

1980 - High water enabled coho to get beyond the C.N.R.culvert. To date no fishway built.

Subject to seasonal fluctuating water levels.

Predators: bears.


Skeena River
Seeley Creek
Seeley Creek
□ - Coho Spawning Grounds
○ - Pink Spawning Grounds
- Railway Bridge (C.N.R.)
- Railway (C.N.R.)
- Road (Highway 16)
- Road Bridge (Highway 16)
N
0
3
kilometres

Sketch of Chicago Creek,
1971.

ESCAPEMENT RECORD FOR CHICAGO CREEK (Seeley Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			400		25	
48			200		25	
49			400		25	
50			400		75	
51			200		25	
52			300			
53			100		75	
54			25		75	
55			200		200	
56			75			
57			75			
58			N/O			
59			N/O			
60			25			
61			75			
62			25			
63			75			
64			75		25	
65			25			
66			75			
67			N/I			
68			N/O			
69			N/O			
70			25			
71			25			
72			N/O			
73			N/O			
74						
75						
76			N/O			
77			N/O			
78			75		200	
79			N/O		25	
80					200	
81					200	
82						
83						
84						
85						

TIMING

ARRIVE			OCT		AUG	
START			E-M OCT		M AUG	
PEAK			M OCT -M NOV		L AUG -E SEP	
END			L OCT - DEC		SEPT	

REMARKS


HAZELTON
PROVINCIAL FOREST
NASS
RANGE
CHIMDEMASH INDIAN RESERVE 2A
Chimdemash Creek
N
0
2
kilometres

NAME OF STREAM CHIMDEMASH CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows W. into Skeena River, E. of S. end of Kitsumkalum Lake, Rge.5, Coast Dist. POSITION 54 128 NE.

LENGTH km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1960 Rocky, steep, swift to Skeena

1960,61,62,65,66. Subject to flash-floods and rapid changes in water levels.

1961 Minor producer--poor spawning area--sparse gravel bars--doubtful fish ever use system--laden with glacial silt--cold mountain runoff. Suggest file be closed.

1962 and 1965 Clear for 1.6 km to canyon.

1962 Trees removed near construction site of new highway 16 bridge.

1964 Silting at mouth where a few pinks spawned in odd years 1950-60

1965 Gravel fan developed at stream mouth. Highway 16 bridge completed.

1967,80 Water course channelized and scoured.

ESCAPEMENT RECORD FOR CHIMDEMASH CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59					N/I	
60					N/O	
61					N/O	
62					N/O	
63					N/O	
64					N/O	
65					N/O	
66					N/O	
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81						
82						
83						
84						
85						

TIMING						
ARRIVE						
START						
PEAK						
END						

REMARKS 1957 Dead Pinks observed near mouth

1967 Report: No fish observed since early 1950-1960



N
Kitsumkalum Lake
MT COUTURE
WESACH MTN
MAROON MTN
MT KHAUSH
MT ALLARD
MT GARLAND
2026
GLACIER PK
1626
MT NIELSON
SLEEPING BEAUTY MTN
MT HENEY
LEAN-TO MTN
MT VANARSDOL
KITSUMKALUM MTN
KITSUMKALUM MOUNTAIN RECREATION AREA
TERRACE
DIST MUN
WATERSHED RESERVE
COPPER MTN
MT LAYTON
NASS RANGES
HAZELTON
SKEENA MOUNTAINS
FOREST
River
Nelson River
Mayo
Sand
750
1500
0
10
kilometres

Clear Creek

For more detail see appendix p. 418

NAME OF STREAM CLEAR CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S. into Kitsumkalum River, Rge.5,Coast Dist.

POSITION 54 128 NW

LENGTH ____ km WIDTH ____ m DRAINAGE ____ km²

COMPOSITION: BEDROCK ____ BOULDER ____ COARSE ____ FINE ____

SILT & SAND ____ UNCLASSIFIED ____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ____ m² SPAWNING AREA ____ m²

DISCHARGE (m³/s) ____

TEMPERATURE (°C) ____

BARRIERS OR POINTS OF DIFFICULT ASCENT

Passable log jams

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- throughout up to 6.4 km
CHINOOK	- up to 6.4 km
COHO	- throughout, especially in main stem headwaters and tributaries.
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS

1948 - small log jams help prevent scouring and cause channel formation.
1949 - average numbers of chinook observed by locals in August.
1950-64 new channel cut for short distance below bridge.
1951 - stream clearance work recommended.
1960,64,67,69,70,71, 1972-78 and 1980 windfalls to 6.4 km
1952 - sockeye runs in August and October.
1960 - live sockeye observed November 25.
1961,64,65,67 very sandy at Kalum Lake head.

Good stream;clear water;large drainage basin,dominated by lake.1961 Opened log-jam at Rosswood to provide spawning salmon with 4.8km of habitat.

Subject to seasonal fluctuation in water levels. Predators: bears and wolves.


Clear Creek
▲ - Chinook Spawning Grounds
▢ - Coho Spawning Grounds
× - Sockeye Sapwning Grounds
Cedar River
Kalum Lake
N
0
5.12
kilometres

Sketch of Clear Creek,1971.

ESCAPEMENT RECORD FOR CLEAR CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947	400	200				
48	750		200			N/O
49	500	OBS	300			
50	1000		200			
51	1000		450			
52	1400		400			
53	500		75			
54	1500		400			
55	* N/O	OBS	N/O			
56	750	200	400			
57	200	400	400			
58	1500	400	400			
59	750	400	N/I			
60	400	200	75			
61	750	400	400			75
62	750	25	200			
63	1500	200				
64	1500	75	1500			
65	750	75	750			
66	750	25	200			
67	500+	100	75			OBS
68	750	75	750			OBS
69	75	75	200			N/O
70	75	75	200			N/O
71	750	200	400			N/O
72	400	75	750			N/O
73	400	200	400			
74	400	200	400			
75	200	200	400			
76	200	75	200			
77	1000	200	400			
78	50	75	200			
79						
80	25	250	400			
81	50	150	400			
82						
83						
84						
85						

TIMING						
ARRIVE						
START						
PEAK						
END						

REMARKS * 1955 Report: Neither Sockeye nor Coho were found on either inspection. This is the first time that this has happened.

N/I - Not inspected

N/O - None observed


SKEENA FOREST
LAKELSE LAKE
Lakelse Lake
LAKELSE LAKE PROV PARK
Lakelse Hot Springs
Catt Pt
Gainey Pt
Squirrel Pt
Mailbox Point
Radio
Campsite
Hatchery Creek
Schulbuckhand Creek
Clearwater Creek
Andalas Cr
Ena Lake
Ena Cr
Onion Lake
Clearwater Lakes
FORCEMAN RIDGE
TREE FARM LICENCE
CANADIAN NATIONAL
24
25
26
27
28
29
30
31
32
2662 T L 8298
2674 T L 8276
2676 T L 11370
2678 T L 11128
2679 T L 11127
2680 T L 11126
2684 T L 8278
2693 T L 8279
2663 T L 11145
2664 T L 11144
2665 T L 11146
6800
6801
6802
6803
6804
6805
6806
6807
6808
6809
5139
3988
3983
5129
5127
5133
5966
6257
6258
6259
6260
6262
6256
N
Clearwater Creek
0
2
kilometres

NAME OF STREAM CLEARWATER CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows N into S. end of Lakelse Lake, Rge.5 Coast Dist.

POSITION 54 128 SW.

LENGTH 6.4km km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- from 1.6km to 4.8km below lake.
CHINOOK	
COHO	- throughout from mouth to headwaters within 2.4 km from lake.
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1960-61 Lakelse data covers this system. Counts taken at Lakelse Fence.
1964 - Large coho assisted through marshy section. Beaver dams stabilized flow.
1965 - Winter logging initiated.
1966 - Logging opened access to upper reaches of stream.
1974 - Some erosion,silting and scouring in lower 6.4km during 2nd week October.
1980 - Numerous fry observed in lower 1.6km of river.

Subject to seasonal fluctuations in water levels.
Predators: bears and eagles.
Beaver dams in lower section removed as required.
First 1.6 km muddy streambed.


Clearwater Lake
Clearwater Creek
Andulus Creek
Scully Creek
Lakelse Lake
ᗐ - Marshy Area
/ - Beaver Dam
▫ - Coho Spawning Grounds
x - Sockeye Spawning Grounds
∴ - Shallow Areas
0
2
kilometres
N

Sketch of Clearwater Creek, 1971.

ESCAPEMENT RECORD FOR CLEARWATER CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			750			
48			400			
49			400			
50			150			
51			450			
52	40		150			
53			75			
54			400			
55			400			
56			400			
57			200			
58			400			
59			N/I			
60			N/I			
61			N/I			
62			N/I			
63			750			
64	204		1500			
65	400		3500			
66	200		750			
67	100		200			
68	400		3500			
69	75		400			
70	75		400			
71	25		200			
72	25		200			
73	25		75			
74	25		75			
75	75		75			
76	25		400			
77	25		75		25	
78	25		400			
79	25		1000			
80						
81	25		350			
82						
83						
84						
85						

TIMING

ARRIVE	AUG		OCT		AUG	
START	M SEPT		L OCT		E SEPT	
PEAK	L SEPT		OCT-M NOV		M SEPT	
END	OCT		L NOV - DEC		L SEPT	

REMARKS N/I - Not inspected


2150
2100
2300
2600
2200
2000
K I S P I O X
3000
3300
3700
2500
2800
3500
3200
1900
2700
2400
1800
1721
Upper Club Creek
1700
Lower Club Creek
Club Lake
S W A N
L A K E
Stephens Lake
R I V E R
N
0 Upper Club Creek 2
kilometres

NAME OF STREAM (Club Creek - Upper)

CONSERVATION DISTRICT 8 **STATISTICAL AREA** 4

LOCATION OF MOUTH Flows E. into Club Lake, Cassiar Dist.

POSITION 55 128 NE

LENGTH 0.8 **km** **WIDTH** **m** **DRAINAGE** km^2

COMPOSITION: BEDROCK BOULDER COARSE **FINE**

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- for 0.4 - 0.8km between Swan and Club Lakes.
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS

1959 - Pan American Oil Company has drilling site in this area and disturbance could have scared away fish.
1961 - Weather conditions inhibited inspection.
1962 - Considerable log debris in stream.
1965 - 1967 Isolated area. Popular sportsmen's paradise.
1978 - Unable to inspect stredm following flood.

Subject to seasonal fluctuations in water levels,but moderated by lakes.
Shallow, clear,stream:rocky streambed,sparse spawning gravel.
Cleared of debris as required.


Upper
Club
Creek
Swan
Lake
Lower
Club
Creek
Kispiox
River
Stephens
Lake
▣ - Coho Spawning Ground
x - Sockeye Spawning
Grounds
N
0
2.56
kilometres

Sketch of Club Creek,(Upper)
1971.

ESCAPEMENT RECORD FOR (Club Creek - Upper)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48	750					
49	400					
50	200					
51	400					
52	200					
53	100					
54	25					
55	200					
56	750					
57	750					
58	750					
59	25					
60	750					
61	N/I					
62	N/O					
63	200		OBS			
64	1500	25	2000			
65	750	25	400			
66	1500		400			
67	2500	100	800			
68	N/O	N/O	N/O			
69	1500	N/O	400			
70	400	N/O	400			
71	1500		400			
72	1500		400			
73	750		200			
74	750		200			
75	750		200			
76	25		400			
77	400		200			
78	N/O		N/I			
79	1500		200			
80	750		250			
81	2000		25			
82						
83						
84						
85						

TIMING

ARRIVE	AUG		AUG			
START	E SEPT	E SEPT	M SEPT			
PEAK	M-L SEPT	M SEPT	L SEPT - OCT			
END	E OCT	L SEPT	OCT - NOV			

REMARKS OBS - Observed
N/O - None observed
N/I - Not inspected

Club Creek (Lower)

See Club Creek (Upper) p.45

NAME OF STREAM (Club Creek-Lower)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows E. into Stephens Lake, Cassiar Dist.

POSITION 55 128 NE.

LENGTH ______ km WIDTH ______ m DRAINAGE ______ km²

COMPOSITION: BEDROCK ______ BOULDER ______ COARSE ______ FINE ______

SILT & SAND ______ UNCLASSIFIED ______

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ m² SPAWNING AREA ______ m²

DISCHARGE (m³/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT ______

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- scattered through length between lakes -- heaviest spawning in 1st .8 km
CHINOOK	
COHO	- scattered through length " " "
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ______

GENERAL REMARKS 1947 - Trout infected with worms in intestines and flesh. Indians reported steelhead run.

1948 - Evidence of egg digging by late spawners.

1954 - Beaver dam .8 km from mouth removed to facilitate upstream migration of sockeye.

1956 - Some egg digging.

1962 - Requires thorough survey.

1965 - Logging development expected.

1966 - Good seeding

1978 - Flood damage minimal because lakes stabilized flow.

1981 - Suggested that baffles be installed in culvert.

Subject to seasonal fluctuations in water levels, but moderated by lakes.

Beaver dams removed as required.


Kispiox River
Swan Lake
Upper Club Creek
Lower Club Creek
Stephens Lake
□ - Coho Spawning Grounds
✕ - Sockeye Spawning Grounds
N
0
2.56
kilometres

Sketch of Club Creek (Lower), 1971.

ESCAPEMENT RECORD FOR (Club Creek-Lower)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947	2500					
48	3500		400			
49	1000		300			
50	500		500			
51	1100		600			
52	3000		500			
53	500		200			
54	200		750			
55	750		400			
56	7500		750			
57	3500		500			
58	3500		1500			
59	750		200			
60	2500		N/O			
61	750		400			
62	OBS					
63	3500		200			
64	1500		1000			
65	750		1500			
66	1500		400			
67	2000		200			
68	3500		1500			
69	3500		750			
70	3500		400			
71	3500		750			
72	3500		750			
73	3500		750			
74	3500		750			
75	3500		750			
76	400		750			
77	3500		200			
78	200		750			
79	3500		400			
80	3000		450			
81	3000		75			
82						
83						
84						
85						

TIMING

ARRIVE	AUG		AUG			
START	L SEPT		M SEPT			
PEAK	M-L SEPT		L SEPT-E OCT			
END	SEPT-OCT		M-L OCT			

REMARKS OBS - Observed


Coldwater Creek
N
0
2
kilometres
R A N G E
M O U N T A I N S
F O R E
S K E E
1000
1200
2661
T L 8280
2676
T L 11370

NAME OF STREAM COLDWATER CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows E. into Lakelse River, W. of Lakelse Lake, Rge. 5, Coast Dist.

POSITION 54 128 SW

LENGTH ______ km WIDTH ______ m DRAINAGE ______ km²

COMPOSITION: BEDROCK ______ BOULDER ______ COARSE ______ FINE ______

SILT & SAND ______ UNCLASSIFIED ______

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ m² SPAWNING AREA ______ m²

DISCHARGE (m³/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT Passable windfalls and log jams at various locations. Recommend that obstruction not be removed as provides protection.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- scattered throughout
COHO	
CHUM	
PINK (ODD YEAR)	- scattered throughout
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ______

GENERAL REMARKS 1980/81 Normal water levels. Some erosion, silting and scouring reported in 1980.

Predators: Eagles and bears.

ESCAPEMENT RECORD FOR COLDWATER CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80		25			565	
81		10			650	
82						
83						
84						
85						

TIMING

ARRIVE		July			Aug.	
START		E. Aug- M.Aug.			M Aug.	
PEAK		Aug-M Sept.			M Aug-E Sep.	
END		Sept.			Sept.	

REMARKS


Comeau Creek
N
0
2
kilometres
SKEENA
RIVER
Comeau
Seeley Lake Provincial Park
Keynton Lake
NEW KITSEGUECLA INDIAN RESERVE 2
Carnaby
16
Yellowhead Highway
CANADIAN NATIONAL
1000
1500
3000
4500

NAME OF STREAM COMEAU CREEK (Carnaby Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows W. into Skeena River, Cassiar Dist.

POSITION 55 127 SW.

LENGTH 1.6 km WIDTH ___ m DRAINAGE ___ km²

COMPOSITION: BEDROCK ___ BOULDER ___ COARSE ___ FINE ___

SILT & SAND ___ UNCLASSIFIED ___

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ___ m² SPAWNING AREA ___ m²

DISCHARGE (m³/s) ___

TEMPERATURE (°C) ___

BARRIERS OR POINTS OF DIFFICULT ASCENT ___

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ___

GENERAL REMARKS 1947 - Reason for decline in spawners unknown.

1951 - Low water delayed run.

1957 - No estimate made.

1958 - Beaver dams present. Small stream used only during big pinks year when water levels favourable.

- Beaver dams caused flooding of adjacent land in 1960,62,66.

- Beaver dams opened in 1961 and 1965

1964 - Swampy,silted. Numerous beaver dams in lower reaches. Steep and turbulent upstream where good spawning gravel found.

1965 - C.N.R.culvert at Woodcock Farm obstructs fish passage.

1966 - Occasionally odd-year flood coho survive to produce minor runs 50-100 Subject to seasonal fluctuations in water levels.

60

ESCAPEMENT RECORD FOR COMEAU CREEK (Carnaby Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48			200		200	
49			200		25	
50			75		25	
51			N/I		200	
52						
53			N/I		300	
54						
55			N/I		N/I	
56			N/I		N/I	
57					Obs.	
58			N/O		N/O	
59			N/I		N/I	
60			N/O		N/O	
61			N/O		N/O	
62			N/O		N/O	
63						
64						
65			25			
66			N/O		N/O	
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81						
82						
83						
84						
85						
TIMING						
ARRIVE						
START			E SEPT-OCT		M AUG	
PEAK			L SEPT-OCT		L AUG	
END			M OCT-NOV		M SEPT	

REMARKS N/I - Not inspected
N/O - None observed
Obs. -Observed


S K E E N A
2500
3300
3000
2800
2400
1900
2000
2200
Carrigan
1700
1700
2000
1800
2200
2100
2500
2500
65
66
68
69
70
71
2400
1600
Cullon
1500
1400
691
690
2200
689
Kline Lake
Creek
3025
1600
3024
2000
3023
683
681
2280
1200
688
427
3026
3020
3020
1700
682
684
694
589
1200
3019
N
Cullon Creek
0
2
kilometres
588
588F
590
1100
2279

NAME OF STREAM CULLON CREEK (Grouse Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S. into Kispiox River, NW of Sammon Lake, Cassiar Dist.

POSITION 55 127 NW

LENGTH 12.8 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Culverts are impassable at low and high velocities unless baffled.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- in upper section and tributaries.
CHUM	
PINK (ODD YEAR)	- in lower reaches
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

good stretch of gravel above culvert increased pink and coho runs.

GENERAL REMARKS

1961 - Windfall debris cleared annually as required.
- Several beaver dams were removed from lower reaches 1962,66,68,70,78,80
- fish stayed below culverts all season.
1964 - Flooding problem.
1965 - Infested with small lamprey. Culverts baffled to reduce velocity.
1974 - 1978 Silting, scouring and erosion in lower reaches.

Subject to seasonal fluctuation in water levels.



Sketch of Cullon Creek 1971.

□ - Coho Spawning Grounds
○ - Pink Spawning Grounds
△ - Chinook spawning grounds
Grouse Creek
Kispiox River
N
0
2
kilometres

64

ESCAPEMENT RECORD FOR CULLON CREEK (Grouse Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			200		200	
48			200		200	
49			400		400	
50			200		400	
51			N/O		N/O	
52					Obs	
53			N/I		N/I	
54			N/I		N/I	
55					Obs.	
56			N/I		N/I	
57			N/I		N/I	
58			N/I		N/I	
59			N/I		N/I	
60			N/I		75	
61			35		400	
62					200	
63			N/I		N/I	
64			400		25	
65			200			
66		25	200			200
67		N/O	25		N/O	
68		25	400		25	
69		25	200		25	
70		25	75		25	
71		25	400		1500	
72		25	75		75	
73		25	75		75	
74		25	75		25	
75		25	75		75	
76		25	75		.25	
77		25	75		75	
78		N/O	75		N/O	
79		N/O	N/O		N/O	
80		N/O	N/O		N/O	
81		N/O	N/O			
82						
83						
84						
85						

TIMING						
ARRIVE		AUG	SEPT		AUG	
START		M AUG	M SEP- M OCT		E -M AUG	MAY
PEAK		L AUG	L SEP- M NOV		M- L AUG	JUNE
END		SEPT	L OCT-DEC		L AUG-L SEP	JUNE

REMARKS N/I - Not inspected
N/O - None observed
Obs.- Observed


Date Creek
N
0
2
kilometres
RIVER
AGWEDIN
INDIAN
RESERVE
Creek
Date
WM
A
R
A
L
397
398
426
2538
2539
2540
2542
2542A
2543
2544
2545
2546
2547
2548
2549
2552
2553

NAME OF STREAM DATE CREEK (Dale Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows E. into Kispiox River, N. of Kispiox, Cassiar Dist.

POSITION 55 127 SW

LENGTH 4.8 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION within 4.8km of mouth.

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- exact grounds unknown.
COHO	- scattered in upper reaches.
CHUM	- in riffles at confluence with Kispiox River.
PINK (ODD YEAR)	- grounds unknown.
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1952,54 Good run of pinks reported by local inhabitants.
1961 - Used in odd years when Kispiox runs are heavy.
1964 - Windfall debris passable. Logging development and improved access expected.
1965 - Logging road expected to extend into area.
1966 - Debris in stream cleared as required.
1967 - Cut bank erosion 0.8km from stream mouth. Flash floods scour gravel bars in lower reaches.
1974 - 20% of lower section affected by scouring, erosion and siltation.
1978 - Possible streambed damage following October flooding. 1979- Scouring and log jams in lower reaches caused by 1978 flood.

Swift, bouldery stream. Basically good gravel. Generally clear.

ESCAPEMENT RECORD FOR DATE CREEK (Dale Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			75		200	
48						
49			400		400	
50			75		200	
51					200	
52					Obs.	
53					200	
54					Obs.	
55			N/I		400	
56			N/I		N/O	
57			N/I		N/O	
58						
59					400	
60			N/O	75		
61			N/I			
62				75		
63						
64		50	300	500		
65				400		
66				200		
67		25		150		
68		25	200			
69			25	25		
70			25			
71			25			
72			25		200	
73			25		75	
74			25		200	
75			25		200	
76			25		75	
77			25		25	
78			25	25	25	
79			N/O			
80						
81			N/O	25		
82						
83						
84						
85						

TIMING						
ARRIVE			E SEP - OCT		AUG	
START			E SEP-M OCT	M - L AUG	L AUG - SEP	
PEAK			L SEP-M NOV	L AUG-E SEP	AUG - SEP	
END			L OCT - DEC	M SEPT	SEP - OCT	

REMARKS N/I - Not inspected
N/O - Not Observed
Obs - Observed


Mount Vanarsdol
WATERSHED RESERVE
Deep Creek
Deep Creek
Dalk ka gila quoeux IR 2
TERRACE DIST MUN
TERRACE MOUNTAIN
Spring Cr
Poultry farm
Sawmill
Butte
Golf Course
CANADIAN
Canyon
Ferry
N
Deep Creek
0
2
kilometres

NAME OF STREAM DEEP CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows SW. into Kitsumkalum River, Rge.5, Coast Dist.

POSITION 54 12m NW

LENGTH 9.6 km WIDTH ______ m DRAINAGE ______ km²

COMPOSITION: BEDROCK ______ BOULDER ______ COARSE ______ FINE ______

SILT & SAND ______ UNCLASSIFIED ______

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ m² SPAWNING AREA ______ m²

DISCHARGE (m³/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT Rock cascade 9.6km upstream from mouth.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- 0.8 to approx.4.8km from mouth.
COHO	- scattered up to 9.6km from mouth
CHUM	- 0.8 to approx 4.8km from mouth
PINK (ODD YEAR)	" " "
PINK (EVEN YEAR)	" " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM Potential of region above cascades not fully assessed in 1965.

GENERAL REMARKS 1951 A few coho fry in L May sampling. 1952 Stream bed silty during Spring sampling period.

1953 - Approx. 180m of stream bed effected by slide triggered by public works road crews.

1958 - Low water restricted entrance of chums and pinks. Coho were observed but not counted.

1960 - Terrace water supply. Considerable muddy sections.

1961 - Logging begun in watershed expected to increase in 1962-63 with proposed sawmill.

1962 - Municipal application to withdraw 22.5 million litres per day protested. Outlet to Kalum River silted for 274m. Some evidence of unspawned pinks and egg digging by late spawning fish.

continued.....

GENERAL REMARKS (cont'd) DEEP CREEK

1962 - Numerous log jams and beaver dams removed. Municipal water supply diversion dam and modification survey pending.
1964 - Terrace municipal drawoff reduces flow considerably and requires study to assure that there are adequate flows during dry years. Protested application approved.
1965 - Suspect that municipal draw down and fish flow requirements cannot be met in years of prolonged drought or if water is impounded above the dam.
1978 - November flood caused some silting and scouring.

Predators: bears and eagles.
Subject to seasonal fluctuations in water levels.
Windfalls, log jams and beaver dams cleared as required.
Good stream, clear water, good gravel, mountain catch basin, deep valley.


Sketch of Deep Creek, 1971.
N
0
.64
kilometres
Deep Creek
- Logging Road
□ - Coho Spawning Grounds
△ - Chinook Spawning Grounds
○ - Pink Spawning Grounds
▲ - Chum Spawning Grounds
- Bridge
Kitsumkalum River

ESCAPEMENT RECORD FOR DEEP CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			75	200	750	25
48		25	75	200	1500	N/O
49		N/O	300	N/O	900	
50			100	200	1200	N/O
51			300	100	900	
52			200	N/O	1500	
53			50	15	600	
54			400	75	750	
55			400	25	750	
56			200	25	750	
57			200	25	1500	
58			Obs	N/O	N/O	
59				25	1500	
60			25	25	25	
61			200	75	7500	
62			200	200	400	
63				75	750	
64		100	1000	75	500	
65		400	1500	75	400	
66		25	400	75	1500	
67		200	400	75	750	
68		75	1500	75	1500	
69		200	400	75	1500	
70		75	400	75	1500	
71		25	400	75	3500	
72		25	400	75	750	
73		25	200	75	400	
74		25	200	25	200	
75		25	200	25	3500	
76		25	200	25	25	
77	25	75	1500	25		200
78		25	200	25	75	
79						
80						
81		N/O	50		400	
82						
83						
84						
85						

TIMING

ARRIVE	AUG	AUG	SEPT	AUG	AUG	
START	E Sept	AUG - E SEP	E SEP -M OCT	M AUG-E SEP	M AUG-E SEP	M MAY
PEAK	M SEPT	M SEPT	M OCT- M NOV	M SEPT	L AUG-M SEP	L MAY
END	L SEPT	L SEPT	OCT - DEC	L SEPT	M - L SEPT	JUNE

REMARKS N/O None observed
Obs Observed


PENINSULA
Woodworth Lake
Mahlon Lake
Bannock Lake
Kergin Lake
Denise Creek
Denise Inlet
Prudhomme Lake
24
23
22
21
20
19
18
17
16
15
14
806
7376
2200
2600
2500
2000
1200
1100
1000
700
500
200
N
Denise Creek
0
2
kilometres

NAME OF STREAM DENISE CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows SW. into Denise Inlet,Rge.5, Coast Dist.

POSITION 54 130 SE.

LENGTH 4.8 km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- within lower 1.6 km
CHUM	" "
PINK (ODD YEAR)	" "
PINK (EVEN YEAR)	" "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1953 Log jam at 3.2 km prevents scouring.

1957 - First 2.0km disturbed by logging operations and littered by debris.

1958 - Logging debris and low flows rendered this stream almost impassable. Recommended that debris be cleared and stream prepared for flash floods.

1962 - Flash flood caused shifting of streambed.

1965 - Sport fishermen reported seeing coho in stream.

1966 - Pink and chum runs have declined steadily since 1950.

Subject to seasonal fluctuations in water levels. Good gravel beds, some bouldersabove 3.2 km. Rapid flow.

Predators: trout, hair seals,bears, eagles and ravens.

Good potential.

continued.....

GENERAL REMARKS (cont'd) DENISE CREEK.

1967 - 5% of first 400m up from mouth affected by siltation and erosion. Flash flood undercut stream banks and caused minor shifts in channels and gravel bars. The reduced pink and chum runs, from this formerly productive stream, might indicate that pollution from the Port Edward pulp mill has had some adverse effect on fry migrating seaward through Wainwright Basin.

1971 - Prior to 1954 this stream supported fair runs of pink and chum salmon. These runs have been almost eliminated and the cause can be directed toward industrial pollution in the area.

1972 - Continued industrial pollution in the vicinity of Morse Basin appears to be adversely affecting this stream.

1973 - Low oxygen levels and toxic pollutants in Morse Basin are probably affecting the productivity of this stream.

1974 - Slight erosion, siltation and scouring. Since 1954 the number of spawners using this stream has dropped to drastically low levels and these runs can be considered nearly extinct.

1976 - Runs in this stream are very low and continue to decline.

1978 - Heavy rains in November may have caused some streambed damage and scouring in the lower reaches.



Denise Creek
To Morse
Basin
Denise
Inlet
To Kloiya
Bay
□ - Coho Spawning Grounds
○ - Pink Spawning Grounds
▲ - Chum Spawning Grounds
N
0
2.0
kilometres

Sketch of Denise Creek,1968.

ESCAPEMENT RECORD FOR DENISE CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947				25	25	
48				3500	3500	
49				1500	4000	
50				300	500	
51			40	100	1000	
52			N/O	2000	3000	
53				1500	750	
54				300	500	
55				N/O	25	
56				14	9	
57			50	1500	400	
58				50	130	
59				25	25	
60			25	N/O	N/O	
61				75		
62				25		
63			NOT INSPECTED			
64			NOT INSPECTED			
65			Obs			
66			400	N/O	N/O	
67			75	25	N/O	
68			N/O	N/O	N/O	
69			75	N/O	N/O	
70			N/O	N/O	N/O	
71			20		6	
72			75	20		
73			20	20	N/O	
74			20	N/O	N/O	
75			20	N/O	N/O	
76			20	20	N/O	
77			20	N/O	N/O	
78			25			
79			N/O			
80			UNK		N/O	
81			25			
82						
83						
84						
85						

TIMING						
ARRIVE			AUG - SEPT	AUG - M SEP		
START			AUG - L SEP	AUG - L SEP	AUG	
PEAK			L SEP- OCT	SEP - E OCT	SEPT	
END			OCT - NOV	M OCT	OCT	

REMARKS N/O - Not inspected
UNK- Unknown


Diana Creek
0
2
kilometres
N
TSIMPSEAN PENINSULA
Diana Lake
Rainbow Lake
Colonel Johnston Lake
Peck Lake
Gamble
Highway
INVERNESS
Control Dam
Rainbow Island Reserve
Dunsmuir

NAME OF STREAM (Diana Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Prudhomme Lake,expansion of Kloiya River,Rge.5,Coast Dist.

POSITION 54 130 SE.

LENGTH 4.8 km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- lower 0.8km in Diana and Prudhomme Creeks.
CHINOOK	
COHO	- lower 1.6km " " "
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1951 - Since dam installed at Prudhomme Lake outlet, the water has backed up in Diana Creek for approx. 0.8km.

1952 - Redd sampled in early March showed good survival rate.

1953 - Dam flooding resulted in lake water backing into stream. Discoloured water in first 1.6km of stream. Flooding provides constant covering of gravel and may become attractive to coho and sockeye spawners now that Kloiya Creek fishway is operating.

1954 - Since Kloiya Creek dam was installed some spawning beds in the lower 0.8km have been lost.

1955,57 High water hampered observation of spawners. Beaver activity has caused numerous channels.

continued

GENERAL REMARKS (cont,d) Diana Creek

1958 - Good seeding resulted from restricted fishing regulations in effect in the Skeena River area.
1960 - A dam is to be constructed in the winter at the outlet of Diana Lake. As a result, water levels in the stream may change.
1961 - Control of Diana Lake Dam by Columbia Cellulose Company has caused drastic fluctuations in water levels. On September 4, 1982 water was completely shut off.
1965 - Some silting resulting from construction on Diana Lake Dam.
1966 - Minimum flow conditions through impounding dam area are 15 cfs mean monthly and 7.5 cfs instantaneous.
1967 - A flood in September caused diversion of channels, stream bank erosion and damaged 25% of the streambed 1.6km from the mouth.
1968 - Because Diana is easily accessible to the public and is a popular recreation area, it requires regular patrols by Fisheries Officers and Guardians.
1972 - Fluctuations in water levels due to impounding dam control.

Predators: eagle,hair seals,bear and wolves.
Good spawning gravel below falls.



Sketch of Diana Creek, 1968.

N
0
1.3
kilometres
To Kloiya Creek
Prudhomme Creek
Prudhomme Lake
Highway 16
Diana Creek
Diana Lake
▭ - Dam and Fish Ladder
▬ - Dam and Spillway
// - Impassable Falls
≈ - Passable Falls
□ - Coho Spawning Grounds
x - Sockeye Sapwning Grounds

ESCAPEMENT RECORD FOR (Diana Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947	75				75	
48	750	25			400	
49	5000				85	
50	5000				N/O	
51	5500					
52	6000					
53	3500		7500			
54	5000		1500			
55	N/R		N/R			
56	2000		3500			
57	750		3500			
58	10000		1500		200	
59	7500		400		N/O	
60	3500		400			
61	3500		1500			
62	3500		3500			
63	7500		750			
64	3500		750			
65	3500	N/O	400		N/O	
66	1500		1500			
67	4500	25	400	N/O		
68	3500	200	1500	25	400	
69	3500	400	1500		N/O	
70	3500	N/O	400		N/O	
71	2000	150	400		N/O	
72	1000	100	350	N/O	N/O	
73	1000	N/O	UNK	N/O		UNK
74	500	N/O	UNK	N/O		
75	1000	N/O	UNK	N/O		
76	700	N/O	UNK	N/O		
77	1500	N/O	N/O	N/O		
78	500		200	N/O		
79	1200		250			
80	*		400			
81	400		200			
82						
83						
84						
85						

TIMING

ARRIVE	AUG - SEPT	AUG - L SEP	M-L SEPT			
START	L AUG -M SEP	SEPT -E OCT	L SEP -E OCT	E SEPT	E SEPT	
PEAK	M SEP-L SEP	SEPT- M OCT	L OCT		M SEPT	
END	M OCT-L OCT	L OCT	L OCT - NOV		L SEPT	

REMARKS 1980 * High water prevented spawning estimates.

N/O - None observed

N/R - No records

UNK - Unknown


Douglas Creek
0
2
kilometres
N
NASS RA
(HAZELTON MOUNT
SKEENA
PROVINCIAL FOREST
WL
Wesach Mountain
6960
6870
6792
CLEAR Creek
Creek
Douglas
Wesach
Creek
14
15
16
17
18
19
20
21
22
23
24
75
76
77
78
79
80
81
82
L 442
L 439
L 438
L 437
L 435
L 436
L 434
L 4368
L 433
L 4367
L 6787
500
900
1000
1500
2000
2500
3500
4000
4500
5000
5500
6000

NAME OF STREAM DOUGLAS CREEK (Dry Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows SW. into Kitsumkalum Lake, Rge.5, Coast. Dist.

POSITION 54 128 NW.

LENGTH 9.6 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1960,76,77 - Windfalls observed between 0.8km and 3.2km.
1962,62 - Cleared windfall debris. 1967,68 Cleared beaver dams.
1962,64 - Some scouring at mouth.
1965,66 - Sandy for 1.6km from swampy bayou to Clear Creek. Improved swampy area near mouth.
1974 - Some erosion and silting during second week in October.
1978 - Beaver dam 3.2km upstream.
1980 - Fish passage hampered by low flow conditions.
Subject to seasonal fluctuations in water levels.
Predators: bears and eagles.
Good spawning and rearing conditions in lower channelled swampy terrain.


Clear
Creek
Kitsumkalum
River
Kitsumkalum Lake
Dry Creek
□ - Coho Spawning Grounds
✗ - Sockeye Spawning Grounds
N
0
5.12
kilometres

Sketch of Dry Creek,

ESCAPEMENT RECORD FOR DOUGLAS CREEK 88 (Dry Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947	75		25			
48						
49	300		100			
50	200		75			
51	200		200			
52	500		400			
53	200		20			
54	200		200			
55	200		75			
56	75		200			
57	25		25			
58	200		200			
59	200		N/I			
60	25		25			
61	200		25			
62	200		25			
63	N/O		N/O			
64						
65	400	OBS	750			
66	25		200			
67	25		25			
68	75		750			
69	200		400			
70	75		400			
71	200		400			
72	200		400			
73	200		200			
74	200		200			
75	200		200			
76	75		200			
77	200		650			
78	75		750			
79						
80	75		225			
81	75		150			
82						
83						
84						
85						

TIMING

ARRIVE	AUG		SEPT-NOV			
START	L AUG-E SEP		L SEP-E NOV			
PEAK	M SEP		M OCT-M DEC			
END	L SEP-L OCT		M NOV - JAN			

REMARKS N/I - Not inspected

N/O - None observed


N
Ecstall River
0
10
kilometres
TREE FARM LICENCE 1
BLOCK 1
Ecstall
River
Khtada Lake
Khtada
MT HAYWARD 1262
Hayward
Big Falls
Herman
Madeline L.
Mudd
Rapids
Kumealon L.
Baker Inlet
Ahks L.
Northness Pt.
Ksngeal L.
Ormond Pt.
Kxngeal Inlet
G R E N V I L L E
Harrion
Ship Anch.
East Inlet
Lower
Ecstall
Tyee
Balmoral
McDonald
Cuthbert
Davis
Windsor
Aviron
Scotia
Carmichael

NAME OF STREAM ECSTALL RIVER

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows NW. into Skeena River near mouth, Rge.5, Coast Dist.

POSITION 54 129SW

LENGTH 100 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Rapids 40.2km upstream.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- in Johnson L. and between Upper and Lower Ecstall Lakes.
CHINOOK	- above Johnson Creek from 3.2km - 32km
COHO	- throughout
CHUM	- just below Johnson Creek to 56km above confluence.
PINK (ODD YEAR)	- scattered throughout, same area as chinooks.
PINK (EVEN YEAR)	- scattered throughout " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1956 - High silt concentration particularly in warm weather.

1963 - High water in L. Sept. hampered inspection.

1965 - Large coho run reported though none observed.

1971 - Silting scouring and erosion in lower reaches (also 72,74,75,76)

1972 - River system is not producing to its capacity with regard to chinook and chums.

1973 - Equipment and manpower restrictions prevented adequate escapement studies through on-site inspection. (also 72,74,76)

1981 - Poor stock assessment due to lack of funds for helicopter survey.

continued.....

ECSTALL RIVER

Large river in remote inaccessible country.

Good looking above tidal waters. Silty up to 43.5km from mouth. Good gravel for 12.9km then 9.6km of sandy silty bottom. Rocky further upstream. Good water supply.

Subject to seasonal fluctuations in water levels.
Predators: hair seals, bears, wolves and wild fowl.
Good spawning conditions and potential.
Spawning areas separated by silty patches.



Sketch of Ecstall River, 1968.

N
0
4.8
kilometres
Beaver Creek
Big Falls Creek
Gib-
ralter
Creek
To Skeena River
Lockerby Creek
Ecstall River
✗ - Sockeye Spawning Grounds
□ - Coho Spawning Grounds
○ - Pink Spawning Grounds
▲ - Chum Spawning Grounds
△ - Chinook Spawning Grounds
Muddy Creek
Clearwater Creek
Tidal Influence
Johnson Lake
Johnson Creek
Rapids
See Inset
Upper Limit of Spawning
Inset
Rapids
Ecstall River
Falls
Lower Ecstall Lake
Falls
Upper Ecstall Lake

ESCAPEMENT RECORD FOR ECSTALL RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947				3500	7500	
48				7500	7500	
49				3500	15000	
50				10000	15000	
51				3000	10000	
52				15000	25000	
53				1500	1500	
54				15000	1000	
55				200	7500	
56				N/O	N/O	
57		3500		1000	1000	
58				3000	20000	
59		3500	N/O	N/O	7500	
60		1500	N/I	N/O	N/O	
61		1500	Obs	35000	Obs	
62		1500	N/O	7500	Obs	
63		3500	Obs		3500	
64		1500	3500	N/O	3500	
65		8750	N/O	3500	20000	
66		3500	3500	15000	1500	
67		1500	1500	12000	750	
68		1500	750	7500	400	
69		750	750	7500	1500	
70		750	750	7500	200	
71		450	2000	2000	1500	
72		1000	1000	20500	N/O	
73	250	3500	UNK	15000	500	
74		2500	UNK	6000	500	
75		2000	UNK	5000		
76		1000	UNK	5000	400	
77		2500	N/O	5000	N/O	
78		750	N/O	3500	200	
79		1000	UNK	3000	N/O	
80						
81		1500	UNK	8000	N/O	
82						
83						
84						
85						

TIMING						
ARRIVE		M AUG	E AUG - SEP	AUG	AUG	
START		M AUG-E SEP	M-L SEPT	M-L AUG	M AUG- E SEP	
PEAK		M SEPT	L SEP-OCT	E SEPT	E-M SEPT	
END		L SEP-M OCT	L OCT -NOV	M SEP-OCT	L SEPT-OCT	

REMARKS N/O - None observed

Obs - Observed

N/I - Not inspected


Erlandsen Creek
N
0
2
kilometres
Erlandsen Creek
Creek
Pine Lake
River
MOUNTAINS
Molybdenum Creek
TREE FARM LICENCE NO 1 BLOCK 4
KITSUM
MOUN
RECRE
ZYMAGOTITZ RIVER
Ski Lift
1000
1500
700
500

NAME OF STREAM ERLANDSEN CREEK 96

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S. into Zymagotitz River, S. of Kitsumkalum L., Rge.5 Coast Dist.

POSITION 54 128 NW.

LENGTH ______ km WIDTH ______ m DRAINAGE ______ km²

COMPOSITION: BEDROCK ______ BOULDER ______ COARSE ______ FINE ______

SILT & SAND ______ UNCLASSIFIED ______

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ m² SPAWNING AREA ______ m²

DISCHARGE (m³/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT ______

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- lower three miles
COHO	
CHUM	- " "
PINK (ODD YEAR)	- lower three miles
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ______

GENERAL REMARKS 1980,81 Water levels normal.
No predators observed.
Some sweepers but passable.

ESCAPEMENT RECORD FOR 98 ERLANDSEN CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80		30		6	36	
81		10		20	50	
82						
83						
84						
85						

TIMING						
ARRIVE		AUG		AUG	AUG	
START		M AUG		M AUG	M AUG	
PEAK		L AUG		L AUG	L AUG	
END		SEPT		SEPT	SEPT	

REMARKS


PONDER
BLOCK 4
MT LEIGHTON
MORTON PK
K I T I M A T
River
Kateen
MT HODGKINSON
MT FINLEY
MT WILLIAM BROWN
Exstew
River
MT SODEN
MT CAMPAGNOLO
KEATS PK
Kasiks
River
KASIKS MTN
S K E E N A
R A N G E
FARM LICENCE 1
BLOCK 1
Arden Lake
Arden
River
Salvus
Devils
Gitnadoix
River
F O R E S T
MT BUCEY
Alder
MT JACKMAN
1500
750
N
Exchamsiks River
0
10
kilometres

NAME OF STREAM EXCHAMSIKS RIVER

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows SE. into Skeena River, Rge.5, Coast Dist.

POSITION 54 129 SE.

LENGTH 24.0 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Passable cascades at 8-16km and 24km.

SPAWNING DISTRIBUTION various locations from mouth to approx. 24 km upstream.

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- usually above 9km upstream
COHO	- in deep water near mouth also in upper reaches and tribs.
CHUM	- 8-9km upstream
PINK (ODD YEAR)	- in lower 4 - 6km of stream.
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1948 - Poor visibility prevented estimate of pink spawners this year and 1949-55 and 1957.

1958 - Indian trappers reported good coho run in November.

1960 - Incomplete coho inspection.

1962 - Parksite popular for late fall coho sport fishing.

- Predators: seals and bears.
- Some erosion and silting in lower areas -- scouring in upper reaches.
- Glacial stream, observation hampered by turbid water.
- Large drainage area, steep glacial runoff approx 40.2km.
- Best observation from helicopter, sandy in lower 4.8km.
- Wide flood plain, swift side channels with gravel bars 30-45m wide.
- Scrub brush and meadows to glaciers, very turbulent in flood.


Exchamsiks River
□ - Coho Spawning Grounds
○ - Pink Spawning Grounds
▲ - Chum Spawning Grounds
△ - Chinook Spawning Grounds
Skeena River
N
0
10.4
kilometres

Sketch of Exchamsiks River, 1971.

ESCAPEMENT RECORD FOR EXCHAMSIKS RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48			50			
49			1000		OBS	
50			300		OBS	
51			800		OBS	
52			1500		OBS	
53			500		OBS	
54			750	OBS	OBS	
55			750	OBS	OBS	
56			1500			
57			400	OBS	OBS	
58			OBS		OBS	
59			NOT INSPECTED			
60			N/O			
61					75	
62			25		25	
63			N/O		N/O	
64		300	400	250	25	
65		75	3500	25	7500	
66		25	200	25	200	
67	400	200	750	25	400	
68		75	750		750	
69		75	750	75	750	
70		75	1500	75	750	
71		75	1500	75	750	
72		75	750	200	1500	
73		75	750	200	750	
74		75	750	200	200	
75		75	750	200	200	
76		75	1500	200	75	
77		75	750	200	400	
78		25	200	50	75	
79		25	500		25	
80						
81						
82						
83						
84						
85						

TIMING

ARRIVE		AUG	SEPT	AUG	AUG	
START		E AUG	M OCT	E AUG	E AUG	
PEAK		M AUG-SEPT	OCT-M NOV	M AUG-SEPT	M AUG	
END		SEPT	OCT-DEC	SEPT	SEPT	

REMARKS N/O - None observed


KITIMAT
RANG
MOUNT ZBURA
MOUNT CONROY
MOUNT MORRIS
MORIN PEAK
Exstew
River
Hudson's Bay Flats
CANADIAN
Clay Creek
Nelson
16
IR 84
IR 83
N
Exstew River
0
10
kilometres

NAME OF STREAM EXTEW RIVER

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows SE. into Skeena River, NE. of Extew, Rge. 5 Coast Dist.

POSITION 54 129 SE

LENGTH 27.4 km WIDTH ______ m DRAINAGE ______ km²

COMPOSITION: BEDROCK ______ BOULDER ______ COARSE ______ FINE ______

SILT & SAND ______ UNCLASSIFIED ______

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ m² SPAWNING AREA ______ m²

DISCHARGE (m³/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT ______

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- 9.6km upstream
CHINOOK	- from 8.0 to 27.4km from mouth
COHO	- upper reaches
CHUM	
PINK (ODD YEAR)	- about 3.2km above bridge
PINK (EVEN YEAR)	- " "
STEELHEAD	- lower reaches

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ______

GENERAL REMARKS 1948 - Poor weather conditions prevented inspection in most years 1948-66

1957 - No damage resulted from gravel removal in Extew Slough.

1964 - Road maintainance work to protect wooden bridge structure.

1965 - Some scouring in lower reaches also in 1968,69. Complete survey of valley required.

1981 - Heavy erosion and silting in lower 6.4km and scouring in upper portion in most years. Alteration in channel.

- Numerous passable log jams along water course.

Subject to seasonal fluctuations in water levels.

Predators : seals and bears.


▢ - Coho Spawning Grounds
Exstew River
Skeena River

Sketch of Extew River, 1971.

N
0
10.4
kilometres

ESCAPEMENT RECORD FOR EXTEW RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48			OBS		OBS	
49			OBS			
50			N/O		OBS	
51			500			
52			OBS			
53			OBS		OBS	
54			OBS			
55						
56					OBS	
57			OBS			
58						
59			NOT INSPECTED			
60			N/O			
61			N/O			
62					200	
63			NOT INSPECTED			
64			N/O			
65				25	750	
66			N/O	N/O	N/O	
67	200		750			
68			75			
69			400			
70			3500			
71			3500			
72		25	3500		1500	
73		200	1500		1500	
74		200	1500		200	
75		200	750		200	
76		75	750		25	
77		75	1500		75	
78		25	750		75	
79		25	700		25	
80						
81						
82						
83						
84						
85						

TIMING

ARRIVE	E OCT	AUG	OCT	AUG	AUG	
START	E OCT	E AUG	NOV	E AUG	E AUG	
PEAK	M OCT	L AUG	M NOV	L AUG	L AUG	
END	L OCT	SEPT	DEC	L AUG	E SEPT	

REMARKS N/O - None observed

(Extew Slough)

See Andeside Creek P.13

NAME OF STREAM (Extew Slough)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows into Skeena River at confluence of Anderside Creek.

POSITION 54 129SE.

LENGTH km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	- primarily in 1.6km of slough and near confluence with
PINK (ODD YEAR)	- " " " Anderside Creek.
PINK (EVEN YEAR)	- " " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1955 - High Silty water prevented accurate observations.

1957 - Indian Trappers reported fair run of coho,also in 1958.

1958 - Turbid water during inspections. Fish seldom enter Andeside Creek from Slough.

1960 - 50% of channel affected by mud and silt from Skeena River.

1961 - Skeena River fallfloods carried considerable sand and silt into slough. Mortalities expected, also in 1965.

1962 - Runs depend of Skeena River flow regime for continued survival.

1964 - Silty bottom deterred spawning. Area used as booming ground without harmful effects.

1966 - Slow winter flows caused drying gravel bars.

- 50% of slough affected by siltation during high water.
- Predation: eagles and bears.

ESCAPEMENT RECORD FOR EXTEW SLOUGH 110

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48				825		
49				30		
50				300		
51				350		
52				50		
53				100		
54				400		
55			OBS	200	200	
56				75		
57				200	75	
58			OBS	400	25	
59				200	75	
60				N/O	25	
61					100000	
62					2500	
63				NOT INSPECTED		
64				N/O	N/O	
65				25	400	
66				30	400	
67				75	200	
68					750	
69				75	750	
70				75	750	
71						
72						
73						
74						
75						
76				25	25	
77				25	25	
78				200	25	
79						
80						
81				20	200	
82						
83						
84						
85						

TIMING						
ARRIVE				AUG	AUG	
START				AUG -E SEPT	E SEPT	
PEAK				M SEPT	M SEPT	
END				L SEPT-OCT	L SEPT	

REMARKS N/O - None Observed


Williams Lake
KISPIOX RIVER
FALLS CREEK
C A S S I
KISPIOX
△ 3201
△ 2150
2100
2200
2300
2400
2600
2700
2800
2900
3000
3200
3300
3400
3700
2000
1900
N
Falls Creek
0
2
kilometres

NAME OF STREAM (Falls Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Swan Lake.

POSITION 55° 45' 128° 40'

LENGTH 0.8 km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Impassable falls 0.8km from mouth.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- to falls or within 1st 0.5km
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM No potential obove falls --inadequate gravel.

GENERAL REMARKS 1948 - Only a short stream and badly overcrowded.
1950 - Beaver dam partially cleared.
1954,55,56,57 Coho schooled of stream mouth--large beaver dam demoved to facilitate migration.
1964 -1966 Recommend complete survey of upper stream area.

Subject to seasonal fluctuation in water levels.
Predators: Bears.


X - Sockeye Spawning Grounds
Falls Creek
Swan Lake
N
0
5.12
kilometres

Sketch of Falls Creek, 1971.

ESCAPEMENT RECORD FOR 114 (Falls Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947	900					
48	15000					
49	2500					
50	500					
51	2000					
52	5000		500			
53	500					
54	75		400			
55	750		400			
56	7500		OBS			
57	3500		100			
58	1500					
59	400		nil			
60	100					
61	N/I					
62	N/O					
63	25		N/O			
64	500					
65	400		400			
66	25					
67	N/R					
68	N/O					
69	N/O					
70	N/O					
71	N/O					
72	25					
73	N/O					
74	N/O					
75	N/O					
76	N/O					
77	N/O					
78	N/I					
79	N/O					
80	N/I					
81						
82						
83						
84						
85						
TIMING						
ARRIVE						
START	AUG-E SEPT		SEPT-E OCT			
PEAK	AUG-L SEPT		SEPT-M OCT			
END	L SEPT-E OCT		L OCT			

REMARKS N/O - None observed
N/I - Not inspected
N/R - No record


N
Fiddler Creek
0
2
kilometres
Skeena River
Chig-in IR
L 910
L 6071
L 909
L 5511
L 5072
L 4486
L 4487
L 5512
L 4485
L 5513
L 6325
L 6327
L 6324
L 6328
L 6323
L 6322
L 6326
L 6321
L 6317
L 6318
L 6319
Abandoned Mine
Creek
Fiddler
RANGES
1500
2000
3700
4000
6000

For more detail see appendix p. 420

NAME OF STREAM (Fiddler Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows N.E. into Skeena River, above Devils Elbow.

POSITION 54° 52' 128° 23'

LENGTH 6.4 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

Rock cascades 3.2 - 6.4km from mouth -- passable depending on flows.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- as far upstream as 9.6km
COHO	- " "
CHUM	- lower 1.6km
PINK (ODD YEAR)	- " "
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

16 - 24km of good gravel above cascades plus numerous tributaries suitable for coho.

GENERAL REMARKS 1948-51 Not formally inspected though residents reported medium - good pink and coho runs.

1952-56 - Residents reported good pink run.

1964 - Good trout fishing, coho taken on rods off stream mouth. Residents reported an early chum run.

1965 - Placer gold mining still actively licensed.

1966 - Placer gold mine licenses issued and active approx. 3.2-4.0 km above C.N.R. line.

1974 - Heavy erosion, silting and scouring in lower river probably destroyed pink and chum runs.

1978 - Erosion and scouring caused by November flood.

1980 - Alluvial fan developed at stream mouth.

continued.

Continuation......

GENERAL REMARKS. (Fiddler Creek)

1981 - Streambed channelized by railroad.

Subject to seasonal fluctuations in water levels.
Stream clear with bouldery substrate--canyon in upper reaches--steep gradient.
Fast flows cause scouring during floods. Dry near mouth at low flows.

Predators: bears.


Sketch of Fiddler Creek,
1971.
N
0
4.11
kilometres
▢ - Coho Spawning Grounds
○ - Pink Spawning Grounds
▲ - Chum Spawning Grounds
△ - Chinook Spawning Grounds
┼┼ - C. N. R. Railway Line
Skeena River
Fiddler Creek

ESCAPEMENT RECORD FOR (Fiddler Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			400		750	
48			OBS		OBS	
49			OBS		100	
50			100		500	
51			N/I		OBS	
52			OBS		OBS	
53			N/I		OBS	
54			N/I		OBS	
55			N/I		OBS	
56			N/I		OBS	
57			N/I		750	
58			N/I		N/I	
59			N/I		N/I	
60			N/I		200	
61			25		400	
62					25	
63						
64			75			
65			400			
66		25	200	25	200	
67		75	400		75	
68		200	750	75	1500	
69		200	200	75	75	
70		200	750	75	750	
71		200	750	75	750	
72		25	750	200	1500	
73		200	750	200	750	
74		200	750	200	200	
75		200	750	75	400	
76		25	750	75	75	
77		75	750	75	200	
78		25	400	75	100	
79		25	200	25	25	
80						
81		N/O	75	N/O	25	
82						
83						
84						
85						

TIMING

ARRIVE		E Aug	Sept	Aug	Aug	
START		E-M Aug	Sept-Oct	E-M Aug	M Aug	
PEAK		L Aug	Sept-Nov	M-L Aug	M-L Aug	
END		Sept	Oct-Dec	Sept	Sept	

REMARKS N/I - Not inspected
N/O - None observed


N
Gitnadoix River
0
10
kilometres
SKEENA
ANGES
FOREST
MOUNTAINS
TREE FARM LICENCE 1
TREE FARM LICENCE 41
MT JACKMAN 1937
Exstew
Hudson Bay Flats
Shames
Zymagotitz River
Gitnadoix
Alastair
Wedeene
MT TEMPLE
MT HOLT
MT MADDEN
DAVIDSON PK
ALASTAIR PK
MT CHARLIE
Amsbury

NAME OF STREAM GITNADOIX RIVER 120

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows N. into Skeena River, Rge.,5 Coast Dist.

POSITION 54 129 SE.

LENGTH 27.3 km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- throughout length of stream.
CHUM	- " " "
PINK (ODD YEAR)	- in lower reaches
PINK (EVEN YEAR)	- " "
STEELHEAD	- " "

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1948 Scouring and water course changes reported in a number of years.

1950,52,58 - Inspection too late to estimate pink spawners.

1960 - Incomplete inspection.

1962 - Sandy at stream outlet.

1963 - Not inspected for coho and chums.

1980 - High incidence of sport fishing during coho run.

Predation: bears,seals,eagles.

Subject to seasonal fluctuation in water levels.



Skeena River
Gitnadiox River
○ – Pink Spawning Grounds
▫ – Coho Spawning Grounds
▲ – Chum Spawning Grounds
△ – Chinook Spawning Grounds
Kadeen River
Alastair Lake
N
0
5.12
kilometres

Sketch of Gitnadnadoix River, 1971.

ESCAPEMENT RECORD FOR GITNADOIX RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			N/I	750		
48		N/O	7500	400	7500	
49			8000	OBS	OBS	
50			5000	500		
51			6000	OBS	OBS	
52			6000	OBS	OBS	
53			5000			
54			7500	OBS	OBS	
55			7500	OBS	OBS	
56		25	15000		OBS	
57		N/O	15000		OBS	
58			35000		OBS	
59			N/I			
60			N/O			
61		OBS	750		3500	
62			3500		400	
63			N/I	N/I	7500	
64			N/O			
65						
66			7500	200	400	
67		750	7500	200	750	
68		750	15000	400	1500	
69		75	7500	400	750	
70		75	7500	400	1500	
71		75	7500	400	1500	
72		75	3500	3500	7500	
73		400	1500	1500	3500	
74		400	750	750	400	
75		400	400	750	1500	
76		75	7500	1500	200	
77		400	400	1500	1500	
78		25	750	200	100	
79						
80		200	2000	1100	7000	
81		200	7000	200	3500	
82						
83						
84						
85						

TIMING						
ARRIVE		JULY	OCTOBER	JULY	AUGUST	
START		AUGUST	OCTOBER	AUGUST	M AUGUST	
PEAK		M AUGUST	M NOVEMBER	L AUGUST	L AUGUST	
END		SEPTEMBER	DECEMBER	SEPTEMBER	SEPTEMBER	

REMARKS N/O - None observed
N/I - Not inspected


N
Glacier Creek
0
2
kilometres
Mount Garland
6652
Icefield
Maroon
3500
6238
5000
Goat
Creek
3500
6000
5300
4000
64
63
62
61
60
59
58
56
55
4500
5339
5351
Icefield
Glacier Peak
6043
5000
NASS
(HAZELTON
Creek
Glacier
3000
1000
1500
Creek
SKE
FOI
1419
1422
1415
1416
1418
1412
1411
1413
500
800
KALUM
WL
Lean-to

NAME OF STREAM GLACIER CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows W. into Kitsumkalum River, S.of Kitsumkalum Lake,
Rge. 5, Coast Dist. POSITION 54 128 NW.

LENGTH 2.4 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE
SILT & SAND UNCLASSIFIED

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT
Impassable falls 2.4km from mouth.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- to falls
CHINOOK	
COHO	- to falls
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS

1965,78,80,81	Erosion,silting and scouring.
1965,66	Some clean up of windfall debris.
	Good gravel and with minor improvements thought suitable for coho production and rearing.
1978,1980	Stream changed course near mouth.
	Predators: eagles and bears.
	Seasonal fluctuations in water levels.
	Mountain glacier fed stream.


- Coho Spawning Grounds
- Kitsumkalum Lake Road
Glacier Creek
Kitsumkalum River
N
0
5.12
kilometres

Sketch of Glacier Creek, 1971.

ESCAPEMENT RECORD FOR GLACIER CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65			750			
66			200			
67			25			
68	750					
69			200			
70			200			
71			200			
72			200			
73			75			
74			25			
75			25			
76			25			
77			25			
78			25			
79						
80			25			
81			N/O			
82						
83						
84						
85						

TIMING

ARRIVE	OCT		OCT			
START	M OCT		OCT-E NOV			
PEAK	M NOV		M NOV			
END	E DEC		DEC			

REMARKS New file 1965


N
Glen Vowel Creek
0
2
kilometres
AGWEDIN
INDIAN
RESERVE 3
KISPAIOX
INDIAN RESERVE 1
Kispiox
RIVER
Glen Vowel Creek
SIK-E-DAKH
Glen Vowell
INDIAN RESERVE 2
ANLAW
INDIAN
RESERVE 4
SKEENA
Quan-skum-kum-mich-mich
Indian Reserve 4
2531
2530
2532
2534
2529
2535
2533
2543
2541
2541F
2542
2917
2918
2513
2126
2157
696
344
345
782
695
697
3633
2266
2265
2267
2268
505
864
2160
2158
2151
3711A
802
900
3800
3960
83
84
85
39
38
37
36
35
34
33
32
31
30
29
28
27

NAME OF STREAM (Glen Vowell Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows into Skeena River North of Hazelton, Cassiar Dist.

POSITION 55 127 SW.

LENGTH 6.4 km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- scattered over lower 4.0km
CHUM	
PINK (ODD YEAR)	- scattered over lower 4.0km
PINK (EVEN YEAR)	- " " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1956,58,61,62,64,65 Low water.

1965 - Good buildup of coho for minor feeder stream.
1965/67 and 1970-80 Blow down debris in lower reaches. Cleared as required.
1966 - Native Indian villagers gaff and net fish.

Fish can only enter stream in years of high flow.
Subject to seasonal fluctuations in water levels.


□ - Coho Spawning Grounds
Glen Vowel Creek
Skeena River
N
0
3
kilometres

Sketch of Glen Vowell Creek,

ESCAPEMENT RECORD FOR GLEN VOWEL CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49			200		200	
50			75		200	
51			75		75	
52			50		100	
53			25		N/O	
54			N/I		75	
55			75			
56			N/O		N/O	
57			N/O		N/O	
58			N/O		N/O	
59			N/I		N/I	
60			N/O		N/O	
61			N/I		N/O	
62			N/O		N/O	
63					N/O	
64			75			
65			75			
66			75			
67			N/O			
68			200			
69			25			
70			25			
71			25			
72			25		200	
73			25		25	
74			25		25	
75			25		25	
76			25		25	
77			25		25	
78			25		N/O	
79			N/O		N/O	
80			N/O		N/O	
81			N/O		N/O	
82						
83						
84						
85						

TIMING						
ARRIVE			SEPT		AUG	
START			E SEP-E OCT		M-L AUG	
PEAK			L SEP-M NOV		L AUG-E SEP	
END			L OCT-DEC		M SEPT	

REMARKS N/I - Not inspected
N/O - None observed


Goat Creek
N
0
2
kilometres
Maroon Creek
Mount Garland
6652
Icefield
Goat Creek
Glacier Park
Icefield
6043
6238
5597
5739
N A S S R
H A Z E L T O N M
Glacier Creek
KITSUMKALUM PROVINCIAL PARK
66
65
64
63
62
61
60
59
1400
1419
1332
3351
4498
6339
4000
4500
3500
2000
1000
5300
5000
6000
BLOCK

NAME OF STREAM GOAT CREEK 132

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows W. into S. end of Kitsumkalum Lake, Rge.5, Coast Dist.

POSITION 54 128 NW

LENGTH 3.2 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

Impassable falls 3.2km from mouth.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- Lakeshore at North end near Kitsumkalum River.
CHINOOK	- Lower 3.2km.
COHO	- Lower 3.2km.
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS

1962,74,78- Flash flooding accompanied by scouring and silting.
1966,67,68- Windfall debris in lower reaches cleared as required.
1976 - Lower 0.8km changed course during October flood.
1977 - Scouring during October flood.
1978,81 - Channel shifted during November flood.

- Seasonal fluctuation in water levels.
- Predators: bears, eagles and poachers.



Sketch of Goat Creek,1971.

□ - Coho Spawning Grounds
x - Sockeye Spawning Grounds
Kitsumkalum Lake
Goat Creek
Kitsumkalum River
N
0
5.12
kilometres

ESCAPEMENT RECORD FOR GOAT CREEK 134

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48			25			
49			70			
50			50			
51			100			
52			100			
53			25			
54			75			
55			75			
56			75			
57			75			
58			75			
59			N/I			
60			N/O			
61			OBS			
62			25			
63			N/I			
64						
65						
66		25	400			
67	25		25			
68			400			
69			75			
70			200			
71	400		200			
72			75			
73			75			
74			75			
75			75			
76			200			
77			200			
78			200			
79						
80		100	25			
81			N/O			
82						
83						
84						
85						

TIMING						
ARRIVE	Sept	July	Oct			
START	M Sept	July - Aug	EOct - ENov			
PEAK	Sept - E Oct	Aug	LOct - MNov			
END	M Oct	Aug - Sept	MNov - Dec			

REMARKS N/I - Not inspected


Hayward Creek
N
0
kilometres
Hayward
Creek
TREE FARM LICENCE 1 BLOCK 1
Mount Hayward △4140
Hayward Lake
ECKSTALL
RIVER
L 5899
L 5892
L 5891
L 5890
L 5888
L 5889
L 5894
△3794
93
92
91
90
89
88
87
86
85

NAME OF STREAM HAYWARD CREEK (Beaver Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows W. into Ecstall R. near mouth. Rge.5. Coast Dist.

POSITION

LENGTH 8.0 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

Impassable falls 6.4 - 8km upstream from mouth.
Cascade type falls 2.4km from mouth.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	see sketch
CHUM	
PINK (ODD YEAR)	see sketch
PINK (EVEN YEAR)	" "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1953 - Inspection incomplete.
1956 - Run size does not warrant stream improvement.
1962 - Survey suggested to determine the extent of spawning gravel above the falls.
1963 - Suggested feasability study of the possibility of using side channels to circumvent falls.
1968-71 Silted throughout tidal influence -- a distance of 1.6km.
1971 - Light scouring of stream bed.
1971-81 Logging debris and subsequent silting seriously affected stream, especially pink spawning grounds in the lower 1.6km.
1974 - Because of reduced runs suggestion made that escapements be included with Ecstall River figures.

continued.....

continuation.....

GENERAL REMARKS HAYWARD CREEK (Beaver Creek)

1978 - Heavy November rains caused considerable streambed damage.
1980 - Poor visibility during September inspection.
Poor spawning habitat,rocky bottom,few gravel bars suitable for spawning.
Subject to seasonal fluctuations in flow rates.
First 4.0km slow flow,turbid,brackish water.
Predators: bears,eagles and seals.



Sketch of Hayward Creek, 1968.

N
0
3.2
kilometres
Port Essington
Ecstall River
Section of Tidal Influence
Pink and Coho Spawning Area When Present
Falls (Impassable)

ESCAPEMENT RECORD FOR HAYWARD CREEK (Beaver Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50					1,000	
51					2,000	
52					4,000	
53					5,000	
54					300	
55					600	
56			N/O		400	
57					20	
58					1,500	
59					50	
60					1,500	
61					25	
62					N/O	
63			25		25	
64						
65			NOT	INSPECTED		
66					750	
67			200		200	
68			N/O		200	
69			N/O	N/O	3,500	
70			N/O		1,500	
71			N/O		N/O	
72			N/O		UNK	
73			N/O		UNK	
74			N/O		N/O	
75			N/O		N/O	
76			N/O		N/O	
77			N/O		N/O	
78			N/O		N/O	
79			N/O		N/O	
80			N/O		N/O	
81			UNK		N/O	
82			UNK		UNK	
83						
84						
85						

TIMING

ARRIVE						
START					Aug	
PEAK					Sept	
END					Oct	

REMARKS


N
Hazelton Creek
0
2
kilometres
2542
KISPAIOX
INDIAN RESERVE 1
Kispiox
RIVER
696
2157
344
345
782
695
697
SIK-E-DAKH
Glen Vowell
INDIAN RESERVE 2
3633
2266
2265
2158
ANLAW
INDIAN
RESERVE 4
SKEENA
505
2267
3711 A
864
2268
504
503
HAZELTON
502
INDIAN RESERVE 1
Hazelton Creek
1611
1612
1613
452
Bk. A
147
501
Hazelton Airfield
4400
4200
3800
3400
3100
2700
2300
2100
1900
1800
1500
1400
1200
1100
1000
900
800

NAME OF STREAM HAZELTON CREEK (Cedar Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows into Skeena River, near four mile canyon on Kitwanga Rd.

POSITION 55 127 SW

LENGTH km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	- Area between mouth and 1.6 km upstream
PINK (EVEN YEAR)	- " " " " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS

1960,1962 Indian smokehouse at stream mouth. Popular trout fishing site.
1961-1977 Sand build-up near mouth
1965 One or two sockeye and pink sighted at Indian Fishery location.
1967 Scouring from late fall floods.
1972 The pinks that spawned during the current season are the first salmon to utilize Cedar Creek since 1962.
1980 Unstable banks 1 km upstream from mouth.

Subject to seasonal fluctuations in flow rates. Steep rocky bottom.
Predators: Bears.

144

ESCAPEMENT RECORD FOR HAZELTON CREEK (Cedar Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947					75	
48					200	
49					200	
50					200	
51					400	
52					500	
53					300	
54					400	
55					400	
56					75	
57					400	
58					*	
59					400	
60					N/O	
61					N/O	
62					100	
63					N/O	
64					N/O	
65					N/O	
66					N/O	
67					N/O	
68			No fish observed			
69			since 1959			
70						
71						
72					1500	
73					400	
74					25	
75					25	
76					N/O	
77					25	
78						
79						
80						
81						
82						
83						
84						
85						

TIMING						
ARRIVE					AUG	
START					M AUG-E SEPT	
PEAK					L AUG-M SEPT	
END					L SEPT	

REMARKS * 1958 No fish entered this stream due to low water

N/O None observed


N
Herman Creek
0
2
kilometres
Terrace Airfield
Thunderbird
Hai Lake
Herman Lake
Mount Herman
4478
1000
500
Herman Creek
Lakgeen
IR 87
Lakelse
Lakelse IR 25
S K E E N A
F O R E S T
Mullar Bay
LAKELSE LAKE PROVINCIAL PARK
LAKE
Furlong Bay
Catt Pt
Gainey Pt
LAKELSE LAKE PROV PARK
Lakelse Lake
Campsite
Sockeye
Williams
Radio
Mailbox Point
25
2 Lines

NAME OF STREAM (Herman Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S. into Lakelse River, W. of Lakelse Lake, Rge.5 Coast Dist.

POSITION 54 128 SW

LENGTH 2.4 km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- scattered to Herman Lake when unobstructed.
CHUM	
PINK (ODD YEAR)	- lower 0.8km
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1964 - Late coho most likely spawned on top of pink redds.

1964 - Some windfall debris. Assessed and cleared as required (65,66,69,72,73)

1965 - Stream clearance improved spawning conditions.

1974 - Log jams 2.4km from mouth

1974 - Some silting and scouring.

1980 - High water removed beaver dams.

Small stream

Subject to seasonal fluctuations in water levels.

Predators: bear and eagles.


▢ - Coho Spawning Grounds
○ - Pink Spawning Grounds
Herman Creek
Lakelse River
N
0
2
kilometres

Sketch of Herman Creek,1971

ESCAPEMENT RECORD FOR 148 (Herman Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			75			
48			25		1500	
49			50		50	
50			N/O		N/O	
51			50		N/O	
52			100		OBS	
53			25			
54			75		75	
55			75		100	
56			75		75	
57			75		OBS	
58			N/I		OBS	
59			N/I		N/I	
60			N/O		N/I	
61			N/I		N/O	
62			N/O		N/I	
63			400		N/O	
64			1,500		500	
65			200		400	
66			200		75	
67			1,500		N/R	
68			200		N/O	
69			200		200	
70			75		400	
71			75		400	
72			25		25	
73			25		25	
74			25		25	
75			25		25	
76					25	
77			25		25	
78						
79			250		25	
80			25		N/O	
81						
82						
83						
84						
85						

TIMING						
ARRIVE			Sept		Aug	
START			E-M Oct		E Aug-LSept	
PEAK			M Oct - MNov		LAug - EOct	
END			L Nov - Dec		Sept - LOct	

REMARKS N/I - Not inspected


Heavenor Creek
0
2
kilometres
N
HEAVENOR CR.
Creek
Brass Lake
Sunday Lake
2564
2555
2551
2552
2553
2549
2550
2554
2563
2558
2561
2562
2556
2557
2560
2559
2546
2545
2547
2548
2538
2574
2578
2579
1569
1559
1557
1563
298
297
296
295
294
399
399A
398
424
875
3027
3035
3036
47
46
45
44
43
42
41
1000
1200
1300
1400
1500
1600
2100
2400
2600
2800

NAME OF STREAM 150 (Heavenor Creek) (McQueen Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows E. into Kispiox River.

POSITION 55° 25' 127° 43'

LENGTH 3.2 km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Beaver dams 1-6km from mouth -- removed as required.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- from mouth to 3.2km upstream
CHUM	
PINK (ODD YEAR)	- " " " "
PINK (EVEN YEAR)	- " " " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1949,50 - Heavy beaver activity though not considered serious.
1961,62 - Suggest file be closed--unproductive stream.
1964 - L.H.and Logging plan to improve access road in 1965.
1965 - Recommend removal of blowdowns.
1966 - File closed.
1968 - File re-opened as McQueen Creek.
Creek runs through partially swampy flat country.
Subject to drying in the summer and generally erratic flow patterns.
1970 - Oct. 3 men spent 3 days removing 14 beaver dams. Work carried out
under Winter Works Stream Clearance Programme.
1972 - Large beaver dam removed by blasting. 1979/80 Beaver dams removed by
S.E.P.Crew.


Murder Creek
McCully Creek
Kispiox River
□ - Coho spawning areas
Heavenor Creek
N
0
2
kilometres

Sketch map of Heavenor Creek, 1971.

ESCAPEMENT RECORD FOR 152 (Heavenor Creek, McQueen Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			75		75	
48			200		75	
49			200		200	
50			75		200	
51			75		75	
52			N/I		N/I	
53			N/I		N/I	
54			N/I		N/I	
55			N/I		200	
56			N/I		N/I	
57			N/I		N/I	
58			N/O		N/O	
59			N/I		N/I	
60			N/O		N/O	
61			N/O		N/O	
62			N/O		N/O	
63			N/O		N/O	
64			N/O		N/O	
65			N/O		N/O	
66			OBS		N/O	
67			-		-	
68			750		-	
69			200		-	
70			200		-	
71			200		200	
72			200		200	
73			200		-	
74			200		-	
75			200		-	
76			200		-	
77			200		-	
78			400		-	
79			25		-	
80			300		-	
81			200		-	
82						
83						
84						
85						

TIMING						
ARRIVE			Sept - Oct			
START			M Sept-MNov			
PEAK			Oct - M Nov			
END			M Oct - Nov			

REMARKS N/I - Not inspected


Hot Springs Slough
0
2
kilometres
N
LAKELSE
Lakelse Lake
SKEENA
FOREST
Squirrel Pt
Lakelse Hot Springs
Hot Springs Slough
Williams Creek
GRACEMAN RIDGE
TREE FARM LICENCE 41
Clearwater Lakes
Onion Lake
CANADIAN NATIONAL
Clearwater Creek
Andalas Cr
End Lake
24
25
26
27
28
29
30
31
32
2662
5139
6800
6801
6802
6803
6804
6805
6806
6807
6808
6809
2675
2674
2676
2677
2678
2679
2680
2684
2689
2690
2691
2693
2695
2668
2663
5120
2666

NAME OF STREAM 154 (Hot Springs Slough)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows int o Lakelse Lake, through hot springs.

POSITION 54 128SW

LENGTH ______ km WIDTH ______ m DRAINAGE ______ km²

COMPOSITION: BEDROCK ______ BOULDER ______ COARSE ______ FINE ______

SILT & SAND ______ UNCLASSIFIED ______

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ m² SPAWNING AREA ______ m²

DISCHARGE (m³/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT Obstructed by windfalls and beaver dams. Recommend removal as required.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- lower portion of slough
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ______

GENERAL REMARKS First report 1980.

ESCAPEMENT RECORD FOR 156 (Hot Springs Slough)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80			24			
81			20			
82						
83						
84						
85						

TIMING						
ARRIVE			Sept.			
START			E. Oct.			
PEAK			L. Oct.			
END			Nov.			

REMARKS


Humpback Creek
N
0
2
kilometres
PORCHER ISLAND
SPILLER RANGES
KITIMAT RANGE
MALACCA PAS
Alice Island
Grace Island
Hunt Point
Island Point
Hunts Inlet
Hunt Inlet
Porcher Island
Ada Islands
Mason Point
Humpback Bay
Lamb Point
Lamppost Islet
Cruice Rock
Lawyer Islands
Foote Creek
Salt Lagoon
Humpback Creek

NAME OF STREAM HUMPBACK CREEK (Humpback Bay Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows N. into Humpback Bay, N. Side of Porcher Island, Rge.5, Coast Dist.

POSITION 54 130 SE.

LENGTH 4.0 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Cascades 4.0km from mouth -- passability depends on water levels.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	Historically throughout, but following clear cut logging in 1966, spawning restricted to lower 1.2-2.4km
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1948-49 Log jam near mouth, however not affecting upstream migration.

1957 - Many small log jams prevent scouring.

1959 - Large number of fish off creek mouth at time of inspection.

1964 - Rock falls 3.2km from mouth.

1965,68 Logging debris removed.

- Silting, scouring and erosion most years between 1966-81.

1966 - Indiscriminate logging practices caused extensive removal of forest cover and debris now washes into the stream at each freshet.

1967 - Logging off forest cover has affected water conditions. Gravel shifts noticeably after each heavy rain with very low water levels between rains. Flood conditions prevail after each heavy rain. Classical example of a small stream in a logged off watershed. Potential limited until there is some forest regrowth.

continued.....

continuation.

GENERAL REMARKS (Humpback Bay Creek)

1971 - Second growth starting to take hold,and stream system beginning to stabilize.
1972 -Some loose eggs observed. Excellent seeding this year.
1976/77 Impassable log jam 2.4km from mouth. Advise removal as causing stream to re-rout and restricting fish to lower 1.2km.
1978 - One log jam opened. Some scouring, loss of spawn and stream bed damage caused by heavy fall rains.
1980 - Approx. 150 pinks died unspawned in estuary. Reason for mortality unknown.

Subject to rapid fluctuations in water levels.
First 1.6km lowland,brackish marshy habitat.
Predators: seals,eagles,otter and wolves.


Malacca Passage
Hump-back Bay
~ - Limit of Tidal Influence
○ - Pink Spawning Grounds
□ - Coho Spawning Grounds
R - Cascades
Humpback Bay Creek
N
0
1.33
kilometres

*Note- Fair Gravel for 4.8 km. Following logging of the watershed in 1966, stream subject to poor water levels, flooding and erosion.

Sketch of Humpback Bay Creek, 1968.

ESCAPEMENT RECORD FOR HUMPBACK CREEK (Humpback Bay Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947					7,500	
48					15,000	
49					25,000	
50					15,000	
51					2,000	
52					30,000	
53					1,500	
54					7,000	
55					750	
56					5,500	
57					3,500	
58					3,000	
59					3,500	
60					1,500	
61					3,500	
62					3,500	
63					3,500	
64					1,500	
65			25		25	
66			400		3,500	
67			75		200	
68			75		3,500	
69			N/0		200	
70			N/0		3,500	
71			N/0		3,000	
72			75		10,000	
73			UNK		750	
74			UNK		1,500	
75			UNK		4,000	
76			UNK		5,000	
77			UNK		1,500	
78			UNK		1,000	
79			UNK		5,800	
80						
81			UNK		1,000	
82						
83						
84						
85						

TIMING						
ARRIVE					EAug-ESept	
START			M Oct		Aug - MSept	
PEAK			L Oct		M-L Sept	
END			M Nov		LSept - Oct	

REMARKS


Johnston Lake
Johnston
68
69
70
71
72
73
74
75
76
K
VK
N
Johnston Creek
0
2
kilometres

NAME OF STREAM JOHNSTON CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows SW into Ecstall River, Rge.4.Coast District.

POSITION 53 129 NE

LENGTH ______ km WIDTH ______ m DRAINAGE ______ km²

COMPOSITION: BEDROCK ______ BOULDER ______ COARSE ______ FINE ______

SILT & SAND ______ UNCLASSIFIED ______

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ m² SPAWNING AREA ______ m²

DISCHARGE (m³/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT

canyon rapid 1.2km from mouth during flood flows.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- evenly from mouth to Johnston Lake
COHO	- " " "
CHUM	- " " "
PINK (ODD YEAR)	- " " "
PINK (EVEN YEAR)	- " " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ______

GENERAL REMARKS 1949,57 - Coho late.

1958 - Expected to become a popular sports-fishing stream and may require strict control in future.

1971 - Slight silting and scouring.

1972,73 Rugged terrain hampers complete assessment.

1978 - November rains may have caused some streambed damage.

- Stream flow stabilized by Johnston Lake.
- Seasonal fluctuations in water levels.
- Predators:seals, bears, wolves, eagles, ravens and mergansers.


Lakeshore Spawning
Johnston
Creek
Johnston Lake
Lakeshore Spawning
To Skeena River
Ecstall
River
Johnston Creek
Ecstall River
✕ - Sockeye Spawning Grounds
▲ - Chinook Spawning Grounds
□ - Coho Spawning Grounds
▲ - Chum Spawning Grounds
N
0
.64
kilometres

Sketch of Johnston Creek, 1972.

ESCAPEMENT RECORD FOR JOHNSTON CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947		3,500	25		750	
48		1,500	25		200	
49		1,500	25		1,000	
50		2,000	20		500	
51		3,000	1,000		500	
52		3,000	500		4,500	
53		1,500	200			
54		3,500	200	1,000	300	
55		3,500	25	25	75	
56		2,500			200	
57		3,500		1,500	1,500	
58		3,000	1,000	500	1,000	
59		3,500	N/I	N/0	1,500	
60		3,500		200	N/0	
61		3,500		1,500	3,500	
62		3,500				
63		1,500		75	3,500	
64		750		400	N/I	
65		NOT INSPECTED				
66		750	400	200		
67		1,500	N/0	400	N/0	
68		3,500		750		
69		7,500	N/0	N/0	N/0	
70		750	N/0	N/0	N/0	
71		300	UNK	300	600	
72		800	500	500		
73		500	UNK	N/0		
74		300	UNK	N/0		
75		300	UNK	N/0		
76		500		500		
77		200		N/0		
78		500		100		
79		500	N/0	150		
80		400	UNK	200		
81		200	UNK	300	25	
82						
83						
84						
85						

TIMING

ARRIVE		Aug	Aug - Sept		Aug - Sept	
START		LAug - Sept	Sept - Oct	Aug	LAug-MSept	
PEAK		E-M Sept	Sept - Oct	Sept	M-L Sept	
END		Sept - E Oct	Oct - Nov	Oct	LSept - Oct	

REMARKS N/I - Not inspected


Johnston Lake
Creek
K
VK
68
69
70
71
72
73
74
75
Johnston
Indian
Reserve

NAME OF STREAM JOHNSTON LAKE

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Expansion of Johnston Creek, N. of Ecstall River, Rge.4, Coast Dist.

POSITION

LENGTH 3.2 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	Lake
CHINOOK	Lakehead.
COHO	"
CHUM	Lake outlet.
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1957 - Inspection made late in year and may account for no trace of fish.
1960,64,66 - Poor observation conditions -- lake turbid due to glacial runoff stream.
1971 - Appear to be two runs of sockeye in system this year.

- Inlet stream considered too cold for spawning and unstable.
-Lake 3.2km long.
Subject to rapid fluctuations in water levels.
Predators: seals,eagles,otter wolves.
first 1.6km lowland, brakish,marshy habitat.



Sketch of Johnston Lake, 1968

N
0
.75
kilometres
Rocky, Glacial Inlet Stream
Steep Mountainous Sides
Beaver Ponds
Johnston Lake
Steep Mountainous Sides
✗ - Sockeye Spawning Grounds
◻ - Coho Spawning Grounds
▲ - Chinook Spawning Grounds
○ - Pink Spawning Grounds
▲ - Chum Spawning Grounds
Swampy Area
Passable Log Jams
R - Rapids
Boulders
R
Ecstall River

ESCAPEMENT RECORD FOR JOHNSTON LAKE

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947	750					
48	400					
49	750					
50	2,500					
51	2,700					
52	5,000					
53	500					
54	500	25		20		
55	25	25		25		
56	5			100		
57	N/O	N/O		N/O		
58	50	N/O		N/O		
59	UNK					
60	UNK					
61	UNK					
62	N/I					
63	UNK					
64	UNK					
65	2					
66	25	25	75	75		
67	750	25	75	75		
68	750	200				
69	400	N/O	N/O	N/O		
70	3,500	N/O	N/O	N/O		
71	5,000	N/O	UNK	N/O		
72	7,500	N/O	UNK	N/O		
73	3,000					
74	750					
75	1,000					
76	3,000					
77	500					
78	500					
79	N/O					
80	N/O					
81	500					
82						
83						
84						
85						

TIMING

ARRIVE	E Aug					
START	L Aug-M Sept	L Aug	L Aug	L Aug		
PEAK	Sept - EOct	E Sept	Sept	E Sept		
END	Oct - M Oct	L Sept - Oct		MSept -Oct		

REMARKS


S T
M O U N T A
E N A
F O R
80
81
82
83
84
85
86
87
88
Creek
Kadeen
ALASTAIR
LAKE
Alastair Peak
ALASTAIR
1000
1500
1900
2000
2100
2700
4200
4500
4800
4900
5000
5200
5800
N
Kadeen Creek
0
2
kilometres

8

NAME OF STREAM KADEEN CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows N. into Gitnadoix River, Rge.5,Coast Dist.

POSITION 54 129 SE.

LENGTH 4.8 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Bouldery cascades beginning at 4.5km may cause migration delays depending on water levels.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- up to 4.8km from mouth.
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1960,61,62,64 Aircraft inspection.

1964 - Suitable gravel bars for coho and chinook spawning.

1967 - Slight scouring

1978 - Heavy silting during November flood.

Large,turbulent, glacial stream. Subject to seasonal fluctuations in water levels.

Predators: bears and eagles.


Sketch of Kadeen Creek,1971
N
0
2.56
kilometres
□ - Coho Spawning Grounds
Kadeen Creek
Kadeen Creek
Alastair
Lake
Gitnadoix River

ESCAPEMENT RECORD FOR KADEEN CREEK 174

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947	25		N/I	25	25	
48	N/0		N/0	N/0	N/0	
49						
50						
51						
52						
53						
54						
55	N/I		N/I	N/I	N/I	
56			750			
57	N/I		N/I	N/I	N/I	
58			1,500			
59			N/I			
60			N/0			
61			N/0			
62			N/0			
63			N/I			
64			N/0			
65			N/0			
66			N/0			
67			25			
68			400			
69			3,500			
70			3,500			
71			3,500			
72			3,500			
73			1,500			
74			1,500			
75			1,500			
76			1,500			
77			750			
78			750			
79						
80			750			
81			500			
82						
83						
84						
85						

TIMING						
ARRIVE			Sept			
START	Aug		M Oct	Aug	Aug	
PEAK			M Nov			
END			Dec			

REMARKS N/I - Not inspected


TREE FARM LICENCE 1
BLOCK 1
KASIKS MTN
Arden Lake
Kasiks River
Telegraph Point
Kwinitsa
Salvus
Devils
Khtada Lake
MT BUCEY
SKIP MTN
Tyee
BALMORAL PK
S K E
F O R
N
TREE FARM LICENCE 1
Kasiks River
0
10
kilometres

NAME OF STREAM KASIKS CREEK

CONSERVATION DISTRICT 8 **STATISTICAL AREA** 4

LOCATION OF MOUTH Flows SE. into Skeena River, Rge. 5, Coast Dist.

POSITION 54 129 SE.

LENGTH 22.5 **km** **WIDTH** ______ **m** **DRAINAGE** ______ **km²**

COMPOSITION: BEDROCK ______ **BOULDER** ______ **COARSE** ______ **FINE** ______

SILT & SAND ______ **UNCLASSIFIED** ______

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ **m²** **SPAWNING AREA** ______ **m²**

DISCHARGE (m³/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT Impassable cascade type falls 22.5km from mouth.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- 6.4 to 25km from mouth
CHINOOK	- " "
COHO	- main stem and small tributaries.
CHUM	- 6.5 to 25km from mouth
PINK (ODD YEAR)	- " "
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ______

GENERAL REMARKS 1955 - Chum run light;pink run moderate.

1959 - Not formally inspected; however, local people report light coho run.

1960 - Silting in lower river most years between 1960-81

1961 - Winter logging and utilization of sloughs for booming proposed.

1962 - No evidence of logging yet.

1964 - Indian food fishery 150-200 coho;good coho sports fishery;18-27kg chinook seen.

1965 - Wide valley flood plain and subject to severe flood attack.

1974 - October floods resulted in scouring, siltation and erosion in the lower river. The pinks and chinook which spawn in this area could have sustained considerable damage. Predators: seals,bears and eagles.


Kasiks River
□ - Coho Spawning Grounds
△ - Chinook Spawning Grounds
○ - Pink Spawning Grounds
▲ - Chum Spawning Grounds
Skeena River
N
0
5.12
kilometres

Sketch of Kasiks Creek,1971

ESCAPEMENT RECORD FOR KASIKS CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50			600	OBS	OBS	
51			1000	300		
52			1000	200		
53			2000	200		
54			1500	200		
55			1500	OBS	OBS	
56			1500	OBS	OBS	
57			3500	OBS	OBS	
58			3500			
59			N/I			
60		25	UNK			
61			UNK	100		
62			1500			
63			UNK	UNK		
64		400	400	400	200	
65		N/O	750	25	400	
66		25	7500	25	N/O	
67	400	25	1500	200	750	
68		400	3500	400	1500	
69		400	3500	400	750	750
70		400	3500	400	1500	
71		400	3500	400	1500	
72		200	1500	400	1500	
73		400	1500	400	750	
74		400	1500	400	200	
75		200	400	200	750	
76		75	750	200	400	
77		200	750	400	750	
78		75	400	200	200	
79						
80		30	1500	250	50	
81		100	2000	25	300	
82						
83						
84						
85						

TIMING						
ARRIVE		Aug	Sept	Aug	Aug	Sept
START	Aug	M Aug	M Oct	M Aug	M Aug	
PEAK		E Sept	L Oct	L Aug	E Sept	
END		Sept	M-L Nov	Sept	M Sept	

REMARKS N/I - Not inspected


N
Khyex River
0
10
kilometres
Tree Farm Licence
Block
Arden Lake
Kasiks
Kasiks Mtn
Exchamsiks
Quottoon
Tyee
Raspberry Is
Port Essington
Ecstall
Balmoral Pt
Windsor Pt
Aylon
Telegraph Point
Cunningham
Veitch Pt
Guest Pt
De Horsey Island
Government Reserve
Rainbow Lake
Skip Mtn
Prudhomme Lake Park
Peninsula
Lachmach
Tree Farm Licence 1

NAME OF STREAM KHYEX RIVER

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S. into Skeena River, NE. of Haysport Rge.5, Coast Dist.

POSITION 54 129 SW.

LENGTH 45 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Impassable cascade falls 29km from mouth.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- upper reaches.
COHO	- throughout.
CHUM	- upper reaches
PINK (ODD YEAR)	- main stem and tributaries
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1948-50 - Some good gravel beds in stream.
1953 - Few males and small females.
- First few km muddy.
1961 - Chums and pinks observed but too early to make an accurate estimate.
1967,72 Only 8km in 1st 29km has satisfactory gravel. 16 km of good gravel above cascades.
1967 - Main stem has large stretches of silty bottom due to tidal action in the lower river.
1968 - Main river turbid,side creeks clear.
Subject to seasonal fluctuations in water levels. Predators: seals, bears, eagles and wolves.


Impassable rapids for two miles
Whitewater Creek
Khyex River
Arden Creek
Sketch of Khyex River
0
1.28
kilometres
N
Δ - Chinook spawning grounds
o - Pink spawning grounds
- Natural Gas Pipeline
- Trail
- C.N.R. Railway
- Railway bridge
- Highway 16
- Highway bridge
Valley Creek
SKEENA RIVER

ESCAPEMENT RECORD FOR KHYEX RIVER 182

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947				750	1500	
48				200	400	
49				400	20000	
50					7000	
51				700	3000	
52				10000	20000	
53				400	1500	
54				500	3000	
55					7500	
56					400	
57				200	3500	
58			200	N/0	1500	
59		400		N/0	3500	
60		750	N/0	N/0	750	
61		OBS		OBS	1500	
62		OBS			3500	
63			1500		3500	
64		25			400	
65		N/0	750	N/0	60000	
66		25	3500	200	400	
67		25	750	200	7500	
68		N/0	750	N/0	3500	
69		N/0	1500	400	7500	
70			N/0	N/0	200	
71		50	UNK	N/0	10000	
72		30	UNK	400	10000	
73		100	UNK	300	20000	
74			UNK	N/0	5000	
75		N/0	UNK	N/0	5000	
76		30	UNK	400	4500	
77		N/0	UNK	N/0	4000	
78		10	UNK	50	500	
79		N/0	UNK	N/0	300	
80		25	UNK	50	1600	
81		UNK	UNK	UNK	2500	
82						
83						
84						
85						

TIMING

ARRIVE		E. Aug	A	E Aug	Aug - Sept	
START		MAug - ESept	MAug - Sept	M-L Aug-Sep	LAug-ESept	
PEAK		Sept	Oct	M Sept	M Sept	
END		M Sept - Oct	L Oct	Sept - Oct	LSept - Oct	

REMARKS


SKE
KITSEGUECLA
INDIAN
RESERVE 1
Skeena
Crossing
Kitseguecla
BM 745
L 823
L 822
Bk A
Bk B
L 2512
L 2511
L 2510
L 2509
L 2508
L 2387
1000
1500
Juniper
P R O
BULKLEY
Kitseguela Logging
Indian Reserve 3
L 4863
L 2606
L 2928
L 2931
Bk C
Bk A
Bk B
L 2930
L 2932
L 2935
L 2934
L 2936
L 2933
L 2423
2500
Kitseguecla
Deep Creek
HAZE
2000
4000
3000
River
F O R E T
N
Kitseguecla River
0
2
kilometres

NAME OF STREAM KITSEGUECLA RIVER (Kitsegukela River)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows NW. into Skeena River, S. of Skeena Crossing,Cassiar Dist.

POSITION 55 127 SW.

LENGTH 32 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- up to approx 12.8km
COHO	- " "
CHUM	
PINK (ODD YEAR)	- " "
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS Silty conditions prevented close inspection of spawners in a number of years between 1954-66

- 1961 Subject to scouring and erosion--noted in most years between 1961-80
- 1964 Sawmill operating near river.
- 1965 Logging and sawmilling road will improve access for inspection.

Glacial stream, swift flow, bouldery,poor salmon producer.
Subject to seasonal fluctuations in water levels.

ESCAPEMENT RECORD FOR KITSEGUECLA RIVER (Kitsegukela River)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			200		200	
48			200		200	
49			400		200	
50			75		200	
51			200		300	
52			N/I		N/I	
53						
54			OBS		OBS	
55			N/I		N/I	
56			N/I		N/I	
57					OBS	
58			N/I		N/I	
59			N/I		N/I	
60			N/0		N/0	
61			N/0		N/0	
62			N/I		N/I	
63			N/0		N/0	
64			OBS		N/0	
65			750			
66		25	200			
67			400			
68			1500			
69			N/0			
70			N/0			
71						
72			N/0			
73			N/0			
74						
75			N/0			
76						
77						
78		25	200			
79			25			
80		25	25			
81		25	200			OBS
82						
83						
84						
85						

TIMING						
ARRIVE		July - Aug				
START		MJuly - EAug			M Aug	
PEAK		LJuly - LAug			L Aug	
END		MAug - Sept			M Sept	

REMARKS N/I - Not inspected


N
Kitsumkalum River
0
20
kilometres
HAZELTON
MOUNTAINS
Kitsumkalum
Kitsumkalum Lake
Nelson River
Mt Caspey
Alder Pk
Cedarvale
Dorreen
Pacific
Lean-to Mtn
Terrace
Kitselas
Lakelse Lake
Mt Garland
Mt Gray
Buildings
Sand L.
Zymacord
Williams
Exstew
Skeena
Tsimpsean
Greenville
Aiyansh
Lakelse
River
2000
3000
4000
1000
500

NAME OF STREAM KITSUMKALUM RIVER (Lower)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S. into Skeena River, Rge. 5, Coast Dist.

POSITION 54 128 NW.

LENGTH km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- scattered from mouth to Kitsumkalum Lake.
COHO	- " " "
CHUM	- " " "
PINK (ODD YEAR)	- " " "
PINK (EVEN YEAR)	- " " "
STEELHEAD	- " " "

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1953 - Light sports catches of coho and chinook.

1954 - Good catch of coho in sports fishery.

1955 - " chinook "

1960 - Silting and erosion also 61,78. Heavy sports fishery near mouth.

1961 - Previous river work seems to have stabilized flow.

1964 - 18-29kg chinook taken in sports fishery.

1965 - Log driving discontinued on river and recommend side channels and sloughs be reopened to provide additional spawning and rearing habitat.

Stream has high silt content --often difficult to estimate number of spawners present. Predators: bears and eagles. Seasonal fluctuations in water levels.


Sketch of Kitsumkalum River (Lower), 1971.

Kitsumkalum Lake
(Kalum Lake)
Nelson Creek
Goat Creek
Star Creek
Glacier Creek
Alice Creek
Lean-to Creek
Luncheon Creek
Kitsumkalum River
(Kalum River)
□ - Coho Spawning Grounds
○ - Pink Spawning Grounds
▲ - Chum Spawning Grounds
△ - Chinook Spawning Grounds
Deep Creek
Spring
Creek
Skeena
River
N
0
5
kilometres

ESCAPEMENT RECORD FOR KITSUMKALUM RIVER (Lower)

YEAR	SOCKEYE	CHINOOK	190 COHO	CHUM	PINK	STEELHEAD
1947		750	3500	750	7500	
48		OBS	OBS	OBS	OBS	
49						
50		OBS	OBS	OBS		
51		OBS	OBS	OBS		
52						
53		OBS	OBS	OBS	OBS	
54		OBS	OBS		OBS	OBS
55		OBS	OBS	OBS	OBS	OBS
56		OBS	OBS	OBS	OBS	OBS
57		OBS	OBS	OBS	OBS	
58		OBS	OBS	OBS	OBS	
59		OBS	OBS	OBS	OBS	
60		OBS	OBS	OBS	OBS	OBS
61		750		75	15000	
62		750	3500	75	200	light
63		OBS	OBS	OBS	3500	
64		2000	1000	300	400	UNK
65		1500	N/O	25	400	good
66						
67						
68		1500	1500	400		OBS
69		7500	400	400		
70		7500	400	400		
71		7500	400	400		
72		3500	400	1500		
73		5000	400	1500		
74		5000	400	400		
75		3500	400	400		
76		3500	750	400	25	
77		9000	400	1000	10000	
78		8000	1500	300	200	
79						
80		4200	750	1800	500	
81		9300	3000	N/O	50000	
82						
83						
84						
85						

TIMING						
ARRIVE		July	Sept	Aug	E Aug	
START		L Aug	M Oct	Aug	M Aug	
PEAK		M Sept	E Nov	M Aug	L Aug	
END		Oct	Dec	Sept	Sept	

REMARKS	chinook	coho	chum	pink	steelhead
48	medium	medium	medium	heavy	
50	below average	light	medium	heavy	
51	light	medium	average	good	
53	average	light	light	average	
54	average	heavy		average	good
55	good	medium	light	heavy	medium
56	medium	good	light	light	average
57	average	normal	light	normal	
58	average	good	light	light	
59	average	very light	light	fair	
60	average	good	light	normal	good


Upper Kitsumkalum River
0
10
kilometres
N
TREE FARM LICENCE 1
BLOCK 4
Kitsumkalum River
Kitsumkalum Lake
MT. WARNIE
MT. HEDEMA
MT. ZOURA
MT. PONDER
MT. LEIGHTON
MORTON PK.
MT. HAGGEN
MT. ALLARD
MT. NEILSON
MT. TRESTON
MT. WILLIAM BROWN
MT. HODGKINSON
MT. FINLEY
MT. HENNEY
SLEEPING BEAUTY MTN.
MT. MORRIS
KITSUMKALA MTN.
MARDON MTN.
T I M A T
Nelson River
River
750
1500

NAME OF STREAM KITSUMKALUM RIVER (Upper) (Beaver River)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows into Kitsumkalum Lake, N. of Terrace, Rge. 5, Coast Dist.

POSITION

LENGTH 40.2 km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (^{o}C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- see sketch map
CHINOOK	- " "
COHO	- " "
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1950-51 Passable log jam about 2.4km from river mouth.

1954,56 - Poor inspection conditions--high, murky water.

1961 - Some stream bed disturbance also in 1968,69,80,81.

1962 - Main coho producer in Kitsumkalum system.

1966 - Access to stream by extension of Columbia Cellulose logging road.

Blow downs in lower 8km -- wide flood plane.

Seasonal fluctuations in water levels.

Predators: eagles and bears.


▲ - Chinook Spawning Grounds
× - Sockeye Spawning Grounds
□ - Coho Spawning Grounds
- Nass Road
- Passable Log Jams
Upper Kitsumkalum River
Mayo Creek
Kitsumkalum Lake
(Kalum Lake)
N
0
10.4
kilometres

Sketch of Kitsumkalum River, (Upper),1971.

ESCAPEMENT RECORD FOR KITSUMKALUM RIVER --UPPER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48			3500			
49			6000			
50						
51			3000			
52			N/I			
53			N/I			
54			N/O			
55			7500			
56			N/I			
57			N/I			
58			7500			
59			N/I			
60			OBS			
61		25	3500	N/O	N/O	
62			750			
63			OBS			
64			1250			
65		6	N/I			
66			N/I			
67		N/O	N/O	N/O	N/O	
68		N/O	N/O	N/O	N/O	OBS
69	200	400	750		N/O	
70	1500	200	750		N/O	
71	750	200	750			
72	400	200	750			
73	400	200	750			
74	400	400	750			
75	400	400	750			
76	400	400	1250			
77	400	400	1500			
78	200	25	400			
79						
80	200	300	400			
81	150	N/O	300			
82						
83						
84						
85						

TIMING						
ARRIVE	Aug	Aug	Oct			
START	E Sept	E Sept	M Oct			
PEAK	M Sept	M Sept	M Nov			
END	M Oct	Oct	Dec			

REMARKS N/I - Not inspected


Kitsumkalum Lake
N
0
2
kilometres
KITSUMKALUM LAKE
Kitsumkalum Lake
Kitsumkalum River
Wesach Creek
Maroon Creek
Goat Creek
Nelson River
Rosswood
KITSUMKALUM PROVINCIAL PARK
BLOCK 4
L 432
L 431
L 4979
L 4980
L 4981
L 7523
L 5664
L 6742
L 6743
1332
1400
1800
74
73
72
71
70
69
68
67
6
5
4
500
600
700
800
1000
1500
1600
2000
3500
4000

NAME OF STREAM KITSUMKALUM LAKE

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Expansion of Kitsumkalum River, North of Terrace,Rge.5 Coast Dist.

POSITION 54 128 NW

LENGTH km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	NE. end of lake near shore at mouths of creeks and Kalum River
CHINOOK	entrance.
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1962 and 1965 Log towing and booming may be initiated at Northern end of lake.

1980 - Numerous trout and fry sighted. Canoe would be best method of counting in this area.

Sand bars dry out at low lake levels--sand deposit in upper end of lake from upper Kiksumkalum R. gradually covering spawning beds. Subject to seasonal fluctuations in water levels.

ESCAPEMENT RECORD FOR KITSUMKALUM LAKE

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947	750					
48	2000					
49						
50	2000					
51	3000					
52	3000					
53	2000					
54	3500					
55	1500					
56	3500					
57	750					
58	1500					
59	1500					
60	200					
61	400					
62	400					
63	400					
64	300					
65	400					
66	400					
67	750					
68	400					
69	400					
70	400					
71	1500					
72	1500					
73	1500					
74	1500					
75	750					
76	400					
77	750					
78	400					
79						
80	150					
81	100					
82						
83						
84						
85						

TIMING

ARRIVE	Aug - Sept					
START	MAug - MSept					
PEAK	LSept - MOct					
END	Oct - M Nov					

REMARKS


N
Kitwanga River
0
10
kilometres
HAZELTON
MOUNTAINS
SKEENA
FOREST
KITEEN
RANGES
BULKLEY
RANGE
Kitwancool
Woodcock
Shandilla
Weber
McCully
MT WEBER 2001
750
1050
1200
1350
1500
1650
1800
600
900
450

NAME OF STREAM KITWANGA RIVER (Kitwancool River)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S. into Skeena River, W of Kitwanga, Cassiar Dist.

POSITION 55 128 SE.

LENGTH 34 km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- in lake
CHINOOK	- see sketch map, scattered spawning throughout although concentrated
COHO	- " " between mouth and lake.
CHUM	- " "
PINK (ODD YEAR)	- " "
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1949 - Good coho run observed by local population.
1952,54,55,58 - More sockeye than usual taken in Indian food fishery.
1953 - Sports fishermen reported good showing of steelhead.
1957,62 - Eggs dislodged from gravel in lower reaches.
1958 - Beaver dam may require removal to enable fish passage next season.
1959 - Beaver dam washed out.
1960 - Counting fence operated at river mouth.
1962 - Heavy U.S.late fall coho and steelhead sports fishery opening.
Logging opened up in lake area. Indian food fishery near village.
1964,55 - Windfall debris requires annual monitoring and removal if necessary.

continued.....

continuation.....

GENERAL REMARKS KITWANGA RIVER (Kitwancool River)

1964 Flood reopened old channel providing increased spawning habitat.
1965 Columbia Cellulose started logging on second tree farm license.
1966 Improved road access has increased sports fishing pressure on chinooks, coho and steelhead, and expanded Native food fishery.
1974 Heavy scouring in lower portions of river likely caused considerable damage to salmon spawn.
1978 Kitwanga Lake kept high water in check so minimal damage to spawning beds.

Subject to seasonal fluctuations in water levels.

Predators: bears and eagles.



Sketch of Kitwanga River, 1971.

Kitwanga River
Kitwanga Lake
Moonlit Creek
Kitwancool Creek
Kitwanga River
Skeena River
▢ - Coho Spawning Grounds
△ - Chinook Spawning Grounds
○ - Pink Spawning Grounds
▲ - Chum Spawning Grounds
N
0
10.4
kilometres

ESCAPEMENT RECORD FOR KITWANGA RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49		600		1000	70000	
50	OBS	700	OBS	5000	100000	
51		400	OBS	5000	100000	
52				800	150000	
53		400		3000	70000	OBS
54	OBS			3500	75000	N/I
55				1500	100000+	
56	3500	400		3500	35000	OBS
57		400		400	200000	
58		400		400	100000	
59		400		750	250000	
60	400	25	25	25	25000	UNK
61	200	25	200	25	110000	25
62	200	25	400	25	77500	75
63			400		170000	
64	200	N/O	200	25	50000	UNK
65	400	25	75	25	35000	OBS
66	200	25	200	25	70000	UNK
67	N/R	N/R	N/R	N/R	100000	
68	200	25	400	75	150000	
69	N/O	75	400	75	175000	
70	N/O	75	400	75	200000	
71	N/O	75	750	75	235000	OBS
72	N/O	75	750	750	175000	
73	N/O	200	400	400	139000	
74	N/O	200	400	400	20000	
75	N/O	200	400	200	200000	
76		75	400	75	200000	
77	25	200	400	200	200000	
78	75	200	2000	200	300000	
79		25	75		15000	
80	100	25	700	N/O	35000	
81	N/O	N/O	200	N/O	110000	
82						
83						
84						
85						

TIMING

ARRIVE	L July	Aug	Sept	E Aug	Aug	
START	M Aug	M Aug	M Oct	M Aug	M Aug	Aug - Nov
PEAK	LAug-E Sept	L Aug	M Nov	L Aug	L Aug	Sept - Dec
END	M Sept	Sept	Dec	LSept- Oct	Sept	Nov - Mar

REMARKS N/I - Not inspected


N
0
Kleanza Creek
2
kilometres
M O U N T A
B U L K L E Y
Kleanza
Creek
KLEANZA MOUNTAIN
KLEANZA CREEK PROV PARK
Noble Five
50
49
48
47
46
45
44
43
42
1500
2000
2500
3000
3500
4000
4400
4500
5000
5100

NAME OF STREAM KLEANZA CREEK (Gold Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows W. into Skeena River, N of Zymoetz River, Rge.5 Coast Dist.

POSITION 54 125 NW.

LENGTH 29.0 km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Cascade falls approx. 3.2km from mouth. Passable during high water levels only.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- upper and lower sections
COHO	- upper sections
CHUM	- near mouth
PINK (ODD YEAR)	- lower 3.2km
PINK (EVEN YEAR)	- " "
STEELHEAD	- occasionally above falls.

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1952,54 - Local reports of steelhead.

1957 - Fisheries Research Board erected a fence on this stream. Approx 300,000 eggs were collected from pinks trapped at the fence. About 1000 pinks spawned below the fence. A few coho and chums were put through the fence.

1958 Fisheries Research Board trapped 660 pinks from which 350,000 eggs were collected for hatchery incubation. 59 steelhead, 7sockeye, 23 coho and 9 chum went through the counting fence. Additional coho and chum migrated up stream after the fence was removed, but water conditions made estimates impossible.

1959,61 Fish taken by Fisheries Research Board for hatchery.

1961,61 Counting fence in operation. Sport fishery indicated a light to poor coho run. Water too high for accurate estimation of spawners.

continued.....

continuation

GENERAL REMARKS KLEANZA CREEK (Gold Creek)

1961 - About 100 coho observed above fence following its dismantling in late October.
- some erosion noted 1964,67,68-75,78

1962 - Hatchery destroyed by fire and approx 1.5 million pink alevins and fry lost.

1965 - Flood scoured and diverted channel causing drying and freezing of pink spawning grounds.

1966 - Flood scoured lower river sections. High runoff velocities now common. Remedial action required to stabilize stream bed.

1974 - Heavy erosion, silting and scouring likely destroyed most pink salmon spawn in the lower 3.2km.
Subject to seasonal fluctuations. Predators: eagles.



Sketch of Kleanza Creek, 1971.

N
0
5.12
kilometres
Kleanza Creek
Skeena River
o - Pink spawning grounds
□ - Coho spawning grounds

ESCAPEMENT RECORD FOR KLEANZA CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			400	200	750	
48			200	400	7500	N/O
49			400	N/O	2000	
50			75	100	9000	
51			300	100	3000	
52			300	25	10000	OBS
53			70	25	5000	
54			200	75	7500	OBS
55						
56			200	25	750	
57					1000	
58	7		OBS	OBS	660	59
59			N/I	N/I	1500	
60			OBS		600	
61	2		27	37	3000	33
62			N/O	N/O	N/O	
63				400	200	
64	UNK	UNK	25	25	75	
65	N/O	N/O	25	25	7500	
66		25	200	25	25	25
67		25	200	25	25	
68			400	N/O	200	
69			750	N/O	75	
70			750	N/O	200	
71			750	N/O	200	
72			750	N/O	1500	
73			400	N/O	400	
74			400	N/O	200	
75		25	75		200	
76		N/O	75		25	
77		25	25		200	
78		N/O	75		75	
79						
80	2	N/O	25		36	
81	N/O	N/O	100		350	
82						
83						
84						
85						

TIMING						
ARRIVE	Aug	July	Sept	Aug		UNK
START	E Aug	E Aug	M Oct	E Aug	Aug	UNK
PEAK	M Aug	L Aug	M Nov	M Aug	Aug-ESept	UNK
END	L Aug	Sept	L Nov - Dec	E-L Sept	L Sept	UNK

REMARKS N/I - Not inspected


Kloiya River
N
0
2
kilometres
Kergin Lake
Prudhomme Lake
Prudhomme Lake Provincial Park
Denise Creek
Denise Inlet
Kloiya Bay
Taylor Lake
Morse Basin
Yellowhead
Water Pipeline
16
12
13
14
15
16
17
18
19
2000
2600
1100
1500
500
200
700
1300
1000
300
130
7375
506
832
975
976
974
645
646
3038

NAME OF STREAM KLOIYA RIVER (Cloyah Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows N.W. into Kloiya Bay, Morse Basin,Rge.5, Coast Dist.

POSITION 54 130SE.

LENGTH km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Passable falls 1.2km from mouth

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- pass through ladder to lake L June to July
CHINOOK	- 1.6km downstream from dam
COHO	- " " most pass through to Diana and Prudhomme System.
CHUM	- in pool below dam
PINK (ODD YEAR)	- mouth of Cloyah Creek to fishway.
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM Fishway 1.6km from stream mouth enables fish to utilize area above the ladders.

GENERAL REMARKS 1952 - Coho observed using fish ladders.

1953,57 Good coho run and fair sockeye run used fish ladders to ascend to Diana Creek spawning gravel at head of Lake System.

1954 - 5000 Sockeye and 1500 coho ascended the fish ladder and entered Diana Creek.

1958 - Large run of sockeye passed through fishway to Prudhomme Lake spawning grounds.

1960 - Fall impassable at low flows which could be corrected by spilling more water.

continued.....

KLOIYA RIVER (Cloyah Creek)

1961,67 - Good sports fishing.
1962 -Columbia Cellulose built a chain fence to stop migrating sockeye from jumping over the sides of the fishway.
1963,66 -Diverting water over the falls in low flows would ensure fish passages.
1967 -.42 m^3/sec established as minimum flow required. The complete absence of pink and chum in recent years may indicate that migrating fry are adversely affected by minor scouring and shifting of gravel bar below water recording station.
1973 Impounding dams at Prudhomme and Diana Lake stabilize system.
1981 10 adult chinook taken for Troller Association incubation box.
1 diseased female collected for pathological testing.

Seasonal fluctuations in water levels.
Predators: hair seals, bears, wolves,wildfowl.
Flow controlled by fishway and checked periodically be regional staff.
Good lake system beyond this stream.



Sketch of Kloiya River, 1971.

N
0
.64
kilometres
To Kloiya Bay
Kloiya River
Kloiya Dam
Taylor Lake
Prudhomme Lake
Prudhomme Creek
Highway 16
Diana Creek
To Diana Lake Diana Dam & Impassable Falls
~ - Limit of Tidal influence
N - Passible Falls
- Trail
)(- Bridge
□ - Coho Spawning Grounds
▲ - Chinook Spawning Grounds
× - Sockeye Spawning Grounds

ESCAPEMENT RECORD FOR KLOYIA RIVER (Cloyia Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947					1500	
48					1500	
49				3500	4000	
50		200			2500	
51		200			2500	
52		N/0	OBS		2500	
53					400	
54					200	
55		200			750	
56		750			750	
57		750			1500	
58		300			500	
59		750		200	750	
60		400		400	N/0	
61		400	400	75	200	
62		200	200			
63		1500	200		3500	
64		400	200			
65		400	400		N/0	
66		750	750		4	
67		350	75	N/0	2	
68		750	750	75	400	
69		400	400	N/0	N/0	
70		400	1500	N/0	N/0	
71		200	100	N/0		
72		150	300	N/0	N/0	
73		250	* 1500	N/0	N/0	
74		200	x 700	25	N/0	
75		200	x 500			
76		100	x 600		100	
77		300	x 500		N/0	
78		300	N/0	25		
79		400	100		20	
80		400	200		50	
81						
82						
83						
84						
85						

TIMING						
ARRIVE		July	Aug - Sept	July	Aug	
START		Aug - Sept	Aug - MOct	L July	M Aug	
PEAK		Aug - EOct	Sept - Nov	Aug	L Aug-Sept	
END		Sept - MOct	L Oct - Dec	E Sept	M Sept -Oct	

REMARKS

* Taken from Guardian's observations at fishway & streampools.

x Includes Diana & Prudhomme Creeks


Kwinitsa Creek
N
0
2
kilometres
Kwinitsa Creek
Kwinitsa
Mount McLean
L 7638
Yellowhead Highway
Telegraph Point
L 7534
McLean Point
Carnation Bank
Carnation Island
Skeena
Scuttsap IR 11
Scuttsap Indian Reserve 11A
L 5247
Khtahda IR 10
L 698
Imkusiyan I R 6B
L 7550
Kilcuteeen I R 9
L 2274
L 1377
L 1378
Khtada
Davis La
L 74
L 76
L 75
L 608
L 694
L 5248
16
Fore
05
04
03
02
01
07
10
1000
1400
1500
2200
2500
3100
600
400
200
700
1200
62
63
64
65

NAME OF STREAM KWINITSA RIVER

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S. into Skeena River, W of Kitsumkalum River, Rge 5, Coast Dist.

POSITION 54 129 SW

LENGTH km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

Impassable cascade-type falls 3.2 km from mouth.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- from mouth upstream 3.2km
CHUM	
PINK (ODD YEAR)	- " "
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS

1960,61,62,64,66 Sand bars developing at mouth for distance of 0.8km

1971-78 - Moderate to heavy silting in lower 1.6km.

Seasonal fluctuations in water levels.

Slow moving stream of glacial origin--clear of debris--swampy.

Predators: seals and bears.



Sketch of Kwinitsa Creek, 1971.

Kwinitsa Creek
Log jam
Skeena River
- Coho Spawning Grounds
- Pink Spawning Grounds
- Impassable Falls
- Highway 16
- Highway Bridge
- C.N.R. Railway
- C.N.R. Railway Bridge
N
0
0.5
kilometres

ESCAPEMENT RECORD FOR KWINITSA RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49					600	
50					1000	
51					800	
52					1200	
53					500	
54					750	
55					750	
56					400	
57					400	
58					25	
59					N/I	
60					N/0	
61					N/0	
62					N/0	
63					N/0	
64					N/0	
65					N/0	
66					N/0	
67					N/0	
68					N/0	
69					N/0	
70					N/0	
71			75		500	
72			25		750	
73			25		75	
74			25		25	
75			25		75	
76			75		25	
77			25		75	
78			75		25	
79			N/I		N/I	
80						
81			N/I		N/I	
82						
83						
84						
85						

TIMING						
ARRIVE			Oct		Aug	
START			M Oct		Sept	
PEAK			E-M Oct		M Sept	
END			Nov - Dec		LSept-EOct	

REMARKS N/I - Not inspected


Gordon Point
CUNNINGHAM
Dodd Passage
One Tree Pt
Village I
Port Simpson Indian Reserve 1
Bath Point
Port Simpson
Stumaun Bay
Town Hill
L 3976
FINLAYSON ISLAND
FINLAYSON ISLAND
IR 19
Sarah Point
PASSAGE
Hook Point
Duncan Point
Fortune Point
Centre Rock
Pender Rock
Dodd Rock
Sparrowhawk Rock
Flat Top Islands IR 2
Otter Anch
Pearl Harbour
Mist Island IR 2
Lahau Creek
Cultivation Point
Burnt Cliff Island
Belletti Point
Burnt Cliff Islands Indian Reserve 30
South Island
Haycock I
North Passage
Whitecliff I
Shattock Hill
Shattock Point
Georgetown Mills
Georgetown Lake
IR 3
East Base Point
Salmon Bight
TSIMPSEAN PENINSULA
Mount Griffin 1364
Mount Ben 1920
INDIAN RESERVE 2
KITIMAT RANGES
(COAST MOUNTAINS
Leading Peak
100
200
300
400
500
700
1000
1500
N
Lahau Creek
0
2
kilometres

NAME OF STREAM (Lahou Creek,Pearl Harbour Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows into Pearl Harbour, W. Coast of Tsimpsean Peninsular S.E. of Finlayson Rge., Coast Dist. POSITION 54 130 NE

LENGTH ______ km WIDTH ______ m DRAINAGE ______ km²

COMPOSITION: BEDROCK ______ BOULDER ______ COARSE ______ FINE ______

SILT & SAND ______ UNCLASSIFIED ______

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ m² SPAWNING AREA ______ m²

DISCHARGE (m^3/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT ______

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- lower 3.2 - 4.8km
CHUM	
PINK (ODD YEAR)	- lower 3.2 - 4.8km
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ______

GENERAL REMARKS 1952 - Pink redds sampled during winter. Eggs -- 4 live, 5 dead
Alevins -- 141 live.
1953 - Log debris 0.8km from mouth.
1961,63 Early run died unspawned due to low water. Later run spawned successfully.
1965 - Severe draught.
1967 - Logging upper reaches has affected water levels.
1971-75 Slight scouring observed also 78,80.
1971-80 Passable log jams variously located within first 8.0km may cause delays in upstream migration depending on water levels. Advised monitoring and removal as required.

continued.....

continuation.

GENERAL REMARKS (Lahou Creek,Pearl Harbour Creek)

1972 - 2 log jams removed opening up an extra 0.8km of gravel.
1973,74 Poor pink returns--common throughout district--coho spawning but difficult to estimate numbers.
1976 Greater return than expected.
1978 Heavy rains caused streambed damage.
1980 Fish arrived earlier than usual.

Water discoloured and brakish for 0.4km,marshy, fully protected during upstream migration.
Good gravel bads in stream,upper portion logged.
Seasonal fluctuations in water levels.
Predators:mergansers,eagles,wolves and bears.


Sketch of LaHou Creek,1971.
N
0
.50
kilometres
- Coho Spawning Grounds
- Pink Spawning Grounds
- Passable Log Jam
- Impassable Log Jam
Good Spawning Gravel
Lake
LaHou Creek
Pearl Harbour

ESCAPEMENT RECORD FOR (Lahou Creek, Pearl Harbour Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48					15000	
49					100000	
50					50000	
51					6000	
52					50000	
53		observed	10		1500	
54					8000	
55					7500	25
56					10000	
57					200	75
58					2000	OBS
59					3500	
60					1500	
61					7500	
62					7500	
63					3500	
64					7500	
65					7500	
66			3500		20000	
67			25		3500	
68			N/O		15000	
69			750		3500	
70			N/O		3500	
71			N/O		9000	
72			UNK		8500	
73			UNK		3200	
74			UNK		3000	
75			UNK		6000	
76					15000	
77			N/O		7000	
78			UNK		7000	
79						
80			UNK		18000	
81					3500	
82						
83						
84						
85						

TIMING						
ARRIVE					M Aug	
START			MSept-E Oct		E Sept	March
PEAK			E-L Oct		M Sept	April
END			E Nov		Oct	April

REMARKS


TREE FARM LICENCE 1
BLOCK 3
Gas Pipeline
TERRACE
Lakelse River
N
0
10
kilometres

NAME OF STREAM LAKELSE RIVER

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows N.W. into Skeena River, Rge.5, Coast Dist.

POSITION 54 128 SW.

LENGTH 19.3 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	See sketch map
CHINOOK	" "
COHO	" "
CHUM	" "
PINK (ODD YEAR)	" "
PINK (EVEN YEAR)	" "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1948,51 Some loss of pink spawn due to overcrowding and coho spawning in same gravel. 49,50,57,64.

1952 - Redd sampling showed some dead pink eggs and eyed coho eggs.
- Slight silting near counting fence 56,57,58.
- Spring and fall run of steelhead.

1954 - Better than average steelhead run,63.

1957 - Recommend counting fence be dismantled as it is decreasing the current and causing silting.

1959 - Counting fence removed.

continued.....

continuation.....

GENERAL REMARKS LAKELSE RIVER

1960 - Numerous log jams noted in the years between 1960 and 1980. Monitored and cleared as required.
1960 - Heavy sports fishery during Kitsumkalum River closure.
1961 - Large pink spawners seen.Heavy sports fishery for coho during pink run. Many pinks damaged by poor fishing techniques.
1962 - Silting observed 64,65,66,74. Logging roads improved access for sports fishermen.Heavily fished on weekends in September-October. Approx.800 coho taken by rods.
1964 - Logging activity increased along stream in 1965,66.
1965 -Logging operations require monitoring to ensure that stream is not damaged.
1981 - Evidence of pink eggs being dug up by coho.

Seasonal fluctuations in water levels.
Predators :eagles, bears, seals and gulls.


Sketch of Lakelse River, 1971

Zymoegotitz River
Shames Creek
Skeena River
Lakelse River
Herman Creek
Williams Creek
Lakelse Lake
Scully Creek
Clearwater Creek
Andulus Creek
Coldwater Creek
× - Sockeye spawning grounds
□ - Coho spawning grounds
○ - Pink spawning grounds
▲ - Chum spawning grounds
△ - Chinook spawning grounds
〰 - Counting fence
N
0
5
kilometres

ESCAPEMENT RECORD FOR LAKELSE RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			3500	400	15000	
48		25	3500	750	75000	N/O
49		16	20000		100000	
50			10000	300	200000	
51		30	20000	5000	200000	
52			15000	N/O	200000	OBS
53		25	12000	25	150000	
54		25	15000	N/O	100000	OBS
55		75	15000	N/O	100000	
56		200	50000+	N/O	75000	OBS
57		200	15000	N/O	200000	OBS
58		200	35000		150000	
59					185000	
60	6000	200	35000	N/O	111000	OBS
61	5500	75	25000	200	311000	UNK
62	5500	25	3500	25	630000	
63		200	35000	400	500000	
64	20758	200	35000	400	1225000	
65	N/O	200	75000	400	837000	OBS
66		200	35000	75	397000	
67		N/R	15000	N/R	235000	
68		200	35000		1112000	
69		400	15000	200	368000	
70		400	25000	400	500000	
71		200	15000	400	250000	
72		200	7500	750	850000	
73		400	3500	750	500000	
74		400	5000	750	250000	
75		400	1500	400	750000	
76		200	750	400	250000	
77		400	3500	400	300000	
78		75	7500		175000	
79		50	5000		300000	
80						
81		250	1500		700000	
82						
83						
84						
85						

TIMING						
ARRIVE	L June	E Aug	Sept	E Aug	Aug	
START	July	M Aug	M Oct	L Aug	L Aug	Sept - Nov
PEAK	E Aug	L Aug	M Nov	M Sept	M Sept	Oct - Nov
END	M Aug	Sept	Dec	Oct	Oct	Jan - May

REMARKS


NASS RANGES
(HAZELTON MOUNTAINS)
SKEENA
FOREST
Lean-to Creek
0
2
kilometres
N
Lean-to Mountain
Creek
61
60
59
58
56
55
24
25
26
27
28
29
30
1422
1416
1411
1418
1413

NAME OF STREAM LEAN-TO CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows SW into Kitsumkalum River,S. of Kitsumkalum Lake.
Range 5, Coast Dist. POSITION 54 128 NW

LENGTH km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE
SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Falls 3 km from mouth of stream.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- In pool at bridge
COHO	- Throughout for 3 km from mouth
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	- At Kalum Lake Road and up 1.6km.
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1962 - Swift and bouldery in sections.
1964 - Good gravel, usually clear water. Could support pinks and chums.
1965 - Log bridge trestles washed out in Oct-Nov flood. Water level must have risen 2-3m.

Water levels 1962-68 Low/Normal. Seasonal high in Oct. and Nov.
1974 Heavy erosion and silting, scouring early October.


□ - Coho spawning grounds
Δ - Chinook spawning grounds
○ - Pink spawning grounds
Lean-To Creek
Kalum Lake Road
Kalum River
N
0
2.50
kilometres

Sketch of Lean-To Creek, 1971.

ESCAPEMENT RECORD FOR LEAN-TO CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947		25			25	
48						
49						
50						
51						
52			75			
53						
54						
55		N/I				
56		N/I				
57		N/I				
58		N/I				
59		-				
60		-				
61		N/O				
62			200			
63		N/O				
64		25	200	-	200	
65			400			
66			400			
67			25			
68			400			
69			75			
70						
71			400			
72			400			
73			75			
74			75			
75			75			
76			200			
77			75			
78			25			
79			-			
80			14			
81			25			
82						
83						
84						
85						

TIMING						
ARRIVE			Oct			
START		Sept	E Oct		Aug	
PEAK		Sept	MOct - MNov		Aug	
END		Sept	LNov - EDec		Aug	

REMARKS N/I - Not inspected


Lockerby Creek
RIVER
N
0
2
kilometres
60
61
62
63
64
65
66
68
69
75
76
77
1600
3500
4000
4500
2100
4200
3200
2900
3900
4100
4238
2000
1100
1300
300
100
400
Knuk-nolks
IR 67
L 2667

NAME OF STREAM (Lockerby Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows into Ecstall River

POSITION

LENGTH 4.8 km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Impassable rock falls approx. 4.8km from mouth

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- for 1.2 - 3.2km above limit of tidal influence
CHUM	- " " "
PINK (ODD YEAR)	- " " "
PINK (EVEN YEAR)	- " " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM good spawning gravel above falls.

GENERAL REMARKS 1953 - Right hand creek logged for distance of 2.4km from mouth. Logs dumped at stream mouth following spawning but,no noticable damage to streambed.

1955 - 5000 coho ascended this stream in early November according to a report by local logger. Also reported good spawning gravel above falls.

1956 - Stream changed course exposing very coarse gravel.

1957 - No logging this year.

1958 - Recommend that stream be changed back to old course if possible since that is where the best gravel is situated.

1961 - Stream appears to be returning to usual following logging.

1962 - Stream bed shifting course with freshets.

continued.....

continuation.

GENERAL REMARKS (Lockerby Creek)

1974 - Slight scouring 75,76,77
- Recommend that Lockerby Creek be included in Ecstall River escapement.
1981 - Full scale survey recommended.

Very bouldery below falls for 0.8km
Good gravel and clear water reported in 50,69 and 81.
2.4km of good spawning area--potential to support larger runs.
Seasonal fluctuations in water levels.
Predators: bears,wolves,seals and eagles.


To Skeena River
(45 km)
Lockerby Creek
□ - Coho Spawning Grounds
○ - Pink Spawning Grounds
// - Impassable Falls
~ - Limit of Tidal Influence
Ecstall River
N
0
1.28
kilometres

Sketch of Lockerby Creek,
1972.

ESCAPEMENT RECORD FOR (Lockerby Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947					7500	
48					1500	
49					5000	
50			2		3000	
51					7000	
52			4		6000	
53			200		1500	
54					1000	
55					7500	
56			2		5	
57			200		750	
58				20	N/0	
59			N/I	N/0	750	
60			N/I	N/0	N/0	
61					7500	
62					75	
63				25	7500	
64						
65					3500	
66			1500		750	
67			1500	N/0	200	
68			N/0	75	N/0	
69			75	75	200	
70			N/0	N/0	75	
71			UNK	N/0	1500	
72			UNK	N/0	300	
73			UNK		750	
74			UNK		N/0	
75			UNK		400	
76			UNK		N/0	
77			UNK		N/0	
78			UNK		N/0	
79			UNK		N/0	
80			UNK		100	
81			UNK		UNK	
82						
83						
84						
85						

TIMING						
ARRIVE					M Aug	
START			E Sept	M Sept	L Aug	
PEAK			M Sept	M Oct	E Sept	
END			L Sept	L Oct	L Sept	

REMARKS N/I - Not inspected


N
0
Lowrie Creek
2
kilometres
SKEENA
CANADIAN NATIONAL
CHIMDEMASH INDIAN RESERVE
KITSELAS INDIAN RESERVE
KLEANZA CREEK PROV PARK
Kitselas Mountain

NAME OF STREAM LOWRIE CREEK (Lowery Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows E. into Skeena River, NE of Usk.Rge. 5 Coast Dist.

POSITION 54 128 NE

LENGTH 3.2 km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	- lower 0.8 - 3.2km
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS

1948,52 - No official inspections, locals report good pink run and medium coho run.

1954 - Medium pink run reported by local residents.

1955,56,57 Fair runs of pinks reported by local residents.

1961,62 Sandy in lower reaches.

1964 - Steep gradient -- scouring observed following freshets 1965-70 and 72-74

1966 - Skeena Forest Products Co. cleared stream of windfalls.

Seasonal fluctuations in water levels.


o- Pink Spawning Grounds
- C.R.N. Railway Line
- Railway Bridge
Lowrie Creek
Skeena River
Usk
N
0
2.56
kilometres

Sketch of Lowrie Creek, 1971.

ESCAPEMENT RECORD FOR LOWRIE CREEK (Lowery Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			400		750	
48			OBS		OBS	
49						
50						
51						
52			OBS		OBS	
53			N/I		N/I	
54			N/I		OBS	
55					OBS	
56					OBS	
57					OBS	
58					OBS	
59					OBS	
60					N/O	
61					N/O	
62					N/O	
63					N/O	
64					N/O	
66					400	
67					25	
68					400	
69					N/O	
70					N/O	
71					N/O	
72					750	
73					200	
74					25	
75					25	
76					N/O	
77					N/O	
78						
79						
80						
81						
82						
83						
84						
85						

TIMING						
ARRIVE			Sept		Aug	
START					E Sept	
PEAK					M Sept	
END					L Sept	

REMARKS N/I - Not inspected


McCully Creek
N
0
2
kilometres
Kispiox R
McCully Creek
Affleck Lake
Brass Lake
Sunday Lake
50
49
48
47
46
45
44
43
K
S
P

NAME OF STREAM McCULLY CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows E. into Kispiox River, Cassiar District.

POSITION 55 127 SW.

LENGTH 9.6 km WIDTH ______ m DRAINAGE ______ km²

COMPOSITION: BEDROCK ______ BOULDER ______ COARSE ______ FINE ______

SILT & SAND ______ UNCLASSIFIED ______

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ m² SPAWNING AREA ______ m²

DISCHARGE (m³/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT ______

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- from mouth approx 9.6km upstream
COHO	- " " "
CHUM	- near mouth
PINK (ODD YEAR)	- " "
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ______

GENERAL REMARKS 1949 Cottonwood log jam near mouth.

1960 - Sandy near outlet to Kispiox River. Log windfall near mouth prevents scouring noted 61,62.

1961 - Chums observed off stream mouth, but none entered stream.

1965 - Good potential as a Kispiox feeder stream.
Windfall debris cleared. Rechannelled to natural channel.

1967 - Slight silting noted 74,79.
Beaver dams noted.

1968 - Log jams at various distances from mouth in most years between 68-79.

1979 - Stream re-routed around large log jams.

continued.....

continuation.

GENERAL REMARKS McCULLY CREEK

Small minor tributary of Kispiox River used by a few fish in good years.
Stream clearance as required.
Seasonal fluctuations in water levels.


McCully Creek
Kispiox River
□ - Coho Spawning Grounds
△ - Chinook Spawning Grounds
▲ - Chum Spawning Grounds
○ - Pink Spawning Grounds
N
0
.64
kilometres

Sketch of McCully Creek, 1971.

ESCAPEMENT RECORD FOR McCULLY CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49					25	
50					75	
51					200	
52					N/I	
53					N/I	
54					N/I	
55					200	
56					25	
57					OBS	
58					75	
59					200	
60					25	
61					N/O	
62				200	25	
63					N/O	
64					N/O	
65			400	3	4	
66		25	200	75	18	
67			25			
68			400			
69			75	25	75	
70			75	25	75	
71			75	25	75	
72			75	200	25	
73		75	75	75	75	
74		75	75	75	25	
75		25	75	25	75	
76		25	75	25	25	
77		25	25	25	75	
78		N/O		25	75	
79		N/O	N/O	N/O	N/O	
80		N/O	N/O	N/O	N/O	
81		N/O	N/O	N/O	N/O	
82						
83						
84						
85						

TIMING						
ARRIVE		Aug	Oct	M Aug	Aug	
START		E Aug	M Oct	L Aug	M-L Aug	
PEAK		L Aug	LOct - MNov	LAug-ESept	LAug-ESept	
END		Sept	Nov - Dec	M Sept	Sept	

REMARKS N/I - Not inspected



McNeil River

Skip Mountain
McNeil River
Aberdeen Point
Skeena Banks
Tyee Bank
Tyee
Hotspring Pt
Raspberry Bluff
Raspberry Islands
Carthew Pt
Skip Rock
Ecstall Island
Port Essington
PORT ESSINGTON IR
Balmoral Peak
3200
3600
3369
16
L 4412
L 4410
L 4416
L 4409
L 4406
L 4408
L 4405
L 4407
L 5485
L 5103
L 1745
L 5467
L 4417
L 6
L 62
L 45
L 11
L 103
L 7291
L 6449
1000
1500
2000
2500
3000
500
N
0
2
kilometres

NAME OF STREAM McNEIL RIVER (Green River)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows into Skeena River E. of Haysport,Rge. 5, Coast Dist.

POSITION 54 129SW.

LENGTH approx 5 km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT impassable rapids approx 5km above lake

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- in upp er section below lake
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	- lower 1.6km above main tidal influence
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1980 - Some silting in lower sections due to tidal influences.
Water levels normal-high.
1981 - Logging activity increased in 1981,but has not encroached on the river yet and no adverse affects have occured to date.

ESCAPEMENT RECORD FOR McNEIL RIVER (Green River)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80			UNK		200	
81			UNK		UNK	
82						
83						
84						
85						

TIMING						
ARRIVE			October		August	
START			November		Sept.	
PEAK			"		M Sept.	
END			"		Oct.	

REMARKS UNK - Unknown


TSIMPSEAN
KITIMAT
VL
COAST
Tuck Inlet
Mount Morse
Mission Mountain
McNichol Creek
Salt Lake Provincial Park
Pillsbury Cove
Schreiber Point
Vigilant Island
Barrett Point
Bacon Point
RUPERT
HARBOUR
PRINCE RUPERT
3975
1991
443
09
10
11
12
13
14
15
16
17
20
21
22
23
24
25
26
27
28
29
30
31
N
Mc Nichol Creek
0
2
kilometres

NAME OF STREAM McNICHOL CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S. into Melville Arm, Prince Rupert Harbour, Rge.5, Coast District. POSITION 54 130 SE.

LENGTH km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

small rock falls 3.2km from mouth

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	- primarily above chute falls 3.2km from mouth
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1953,57,73 Boulder turbulence 1.6km from mouth. Water very discoloured.

1964 - Not officially inspected;however, sports fishermen reported 500 fish.

1968 - Spawning fish in this stream are molested by local children. Recommend closure during spawning season.

1971 - Silting noted most years between 71 and 80.

1978 - November floods possibly damaged stream bed.

Good gravel bars above falls -- fair spawning stream--low marshy surrounding country.

Seasonal fluctuations in water levels.

Predators: hair seals, eagles, bears.


N
Sketch of McNichol Creek,
1971.
0
.48
kilometres
Neither Fork Inspected
Good Spawning
Gravel
Coarse Boulders
Very Limited Spawning Area
Lake
Lake
McNichol Creek
○ - Pink Spawning Grounds
- - Trail
Z - Passable Falls (Chute Type)
Melville Arm

ESCAPEMENT RECORD FOR McNICHOL CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947					750	
48					7500	
49					4000	
50					5000	
51					500	
52					3000	
53					75	
54					5000	
55					75	
56					40	
57					25	
58					500	
59					N/I	
60					1500	
61					1500	
62					200	
63					750	
64					400	
65					N/O	
66			1500		3500	
67			N/O		200	
68					7500	
69			200		750	
70			200		750	
71			N/O		800	
72			N/O		750	
73			UNK		200	
74			UNK		175	
75			UNK		150	
76			UNK		N/O	
77			UNK		200	
78			UNK		600	
79			UNK		75	
80			UNK		400	
81					400	
82						
83						
84						
85						

TIMING						
ARRIVE					M-L Aug	
START			E Oct		LAug-E Sept	
PEAK			M Oct		E-L Sept	
END			L Oct		L Sept-Oct	

REMARKS N/I - Not inspected


Madeline Creek
0
2
kilometres
N
Madeline Lake
Madeline Creek
Madeline Creek
McKnight Creek
McKnight Cr.
Creek
83
82
81
80
79
78
77
76
75
74
73
72
71
K I T
3379
1000
1500
2000
3000
4000
500
100
1700
1300

NAME OF STREAM MADELINE CREEK (Gibraltar Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows into Ecstall River

POSITION

LENGTH 3.2 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Impassable rock falls 3.2km from mouth

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	evenly distributed in lower reaches above tidal influence.
PINK (EVEN YEAR)	" " " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS approx. 3.2km of good spawning gravel.

1966 - Coho reported by loggers.

1968 - Silted in region of tidal influence also noted 69,70.

1973 - This stream has limited areas suitable for salmon spawning. The lower 0.4km is turbid with mud stream bed and influenced by tides in Ecstall River. Approx 0.4km of questionable spawning ground is situated before the stream gradient. Above this area the stream is a continuous series of cascading rock falls.

1974 Recommend that escapement figures be included in Ecstall River spawning report in future.

continued.....

continuation.....

GENERAL REMARKS	MADELINE CREEK (Gibraltar Creek)
1976,77,79,80	Restricted spawning area. Slight erosion problems.
1978	Heavy rains in November caused considerable stream bed damage.
1980	Generally clear water. Stream inclined to flood and change course. Coarse gravel is not easily disturbed.Good gravel sections. Predators, bears and eagles.



Sketch of Madeline Creek, 1968.

N
0
2.0
kilometres
Big Falls Creek
Ecstall River
o - Pink Spawning Grounds
~ - Limit of Tidal Influence
// - Falls and Cascades
Gibralter Creek
Madeline Lake

ESCAPEMENT RECORD FOR MADELINE CREEK (Gibralter Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57				3500		
58				N/O		
59				N/O		
60				N/O	25	
61					7500	
62					N/O	
63					7500	
64					N/O	
65					7500	
66				OBS	200	
67					200	
68					N/O	
69					400	
70					N/O	
71					400	
72					N/O	
73					N/O	
74					N/O	
75					N/O	
76					N/O	
77					N/O	
78					N/O	
79					N/O	
80					N/O	
81					N/I	
82						
83						
84						
85						

TIMING

ARRIVE					Aug - Sept	
START				Aug	LAug-M Sept	
PEAK				Sept	L Sept	
END				Sept	Sept - Oct	

REMARKS


N
Magar Creek
0
10
kilometres
SKEENA
RANGES
FOREST
TREE FARM LICENCE 1
TREE FARM LICENCE 41
Exstew
Hudson Bay Flats
MT JACKMAN
1937
Gitnadoix River
Magar
Wedeene
Dahl
Alastair Lake
MT TEMPLE
MT CHARLIE
DAVIDSON PK
MT BUCKY
750
1500

NAME OF STREAM MAGAR CREEK (Dog Tag Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows N.W. into Gitnadoix River, Rge. 5, Coast Dist.

POSITION 54°12' 129° 12'

LENGTH 6.4 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- spawning for 1.6 - 6.4km upstream from Gitnadoix River.
COHO	- " " " "
CHUM	- " "
PINK (ODD YEAR)	- " "
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1948,49,55 Lower end of stream covered by fine silt.
1949-64 High water prevented estimate of spawners and species in a number of years.
1957 -Inspection too late to get estimates of spawners.
1964,65,66 Air inspection only.
1971 -Unable to inspect this stream from ground due to large log jam on Gitnadoix River.
1974 -Some erosion,silting and scouring in lower end in mid.October.
1978 - " " " caused by November flood.
Flood conditions in Gitnadoix System limit times that creek can be inspected.
Seasonal fluctuations in water levels. Predators: bears.


Dog Tag
Creek
Gitnadiox
River


N
0
10.4
kilometres

Sketch of Magar Creek,
1971.

ESCAPEMENT RECORD FOR MAGAR CREEK (Dog Tag Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48				750		
49				OBS		
50				600		
51				550		
52				200		
53				OBS	OBS	
54			OBS	OBS	OBS	
55			OBS	OBS	OBS	
56			N/O	N/O	N/O	
57				OBS	OBS	
58				OBS	OBS	
59			NOT INSPECTED			
60			NOT INSPECTED			
61			OBS	OBS	OBS	
62			200		25	
63			NOT INSPECTED			
64			N/O	N/O	N/O	
55			NOT INSPECTED			
66		N/O	400	25	N/O	
67		25	400	25	25	
68		25	400	25	25	
69		N/O	400	N/O	400	
70		25	400	25	400	
71		N/O	N/O	N/O	N/O	
72		25	400	25	400	
73		25	200	25	400	
74		25	750	75	400	
75		25	400	25	400	
76		25	750	25	25	
77		25	400	25	200	
78		25	400	750	25	
79		25	75	2000	25	
80						
81		10	100	200	100	
82						
83						
84						
85						

TIMING						
ARRIVE		July	Sept - Oct	E Aug	E Aug	
START		Aug	Oct - Nov	E Aug	E Aug	
PEAK		M Aug	M Nov	M-L Aug	E-M Sept	
END		Sept	Dec	LAug -Sept	L Sept	

REMARKS


Moore Cove Creek
N
0
2
kilometres
Brown Lake
KITIMAT
(COAST MOUNT
MT ELWIN
Kromann
Creek
Moore Cove Creek
Moore Cove
Fleming Bay
Chail Point
1003
5954
214
213
212
215
211
1000
1500
2000
2300
2500
500
400
200
100
1800

NAME OF STREAM MOORE COVE CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows into Moore Cove, E. shore of Telegraph Passage, Rge.5, Coast Dist. POSITION 54N.130W.

LENGTH 6.4 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

Small rock falls at 2.4km up right fork. Passable at high water levels.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- upper reaches and headwaters
CHUM	- lower reaches
PINK (ODD YEAR)	- scattered throughout
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1949 - Suggested remedial work on falls to make them passable regardless of flow levels.

1950 - Chums distributed over tidal and non-tidal stream for approx. 73m

1953 - Low water periods during the run could block ascent to gravel. Guardian provided bypass by drill and hammer work along side during normal water levels.

1956 - Obstructions on both forks largely eliminated.

1961 - Some loss of early run fish due to low water levels.

1962 - Suggest proper survey be made of spawning area above falls.

1963 - Low water delayed upstream migration until early September.Some fish died before spawning.

continued.....

Continuation.

GENERAL REMARKS
MOORE COVE CREEK

1964 - Suggest access over falls be improved.
1965 - Guardian transported 3600 fish over obstruction during draught period.
1967 - Logging within 0.4km of stream. Recommend close monitoring of situation.
1968 - Falls blasted out to relieve upstream migration at low flows.
1973 - Decline in pinks--reason unknow.Suggest remedial blasting of cascades in lower 3.2km of the river.
1975 - Cascades successfully blasted.
1976 - High flows and water discoloration impaired inspection.
1978 - High flows may have caused some stream bed damage.
1981 - Summer inspection revealed an abundance of coho fry.

Seasonal fluctuations in water levels.
Predators: hair seals, wolves, bears and birds.
Good gravel beds below falls.


Telegraph Passage
○ - Pink Spawning Grounds
□ - Coho Spawning Grounds
N - Passable Cascade-type Falls
~ - Limit of Tidal Influence
■ - Guardian's Cabin
Moore Cove
Moore Cove Creek
Lake
N
0
1.6
kilometres

Sketch of Moore Cove Creek.

ESCAPEMENT RECORD FOR MOORE COVE CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			1500		7500	
48			750		35000	
49			100		50000	
50			400	500	10000	
51			1000		40000	
52			N/O		50000	
53					15000	
54			N/O		15000	
55			250		15000	
56					15000	
57					75000	
58					75000	
59					60000	
60			OBS		35000	
61			3500		35000	
62			OBS		100000+	
63			OBS		75000	
64			OBS		35000	
65					7500	
66			7500		35000	
67			400		10000	
68			3500		130000	
69			3500		15000	
70			750		35000	
71			900		65000	
72			500		50000	
73			700		3700	UNK
74			600		40000	
75			UNK		30000	
76			UNK		20000	
77						
78			UNK		10000	
79			2000		6000	
80			UNK		8500	
81			UNK			
82						
83						
84						
85						

TIMING

ARRIVE			MSept - EOct		LAug-MSept	
START			MSept - EOct		ESept-LSept	
PEAK			LSept - MOct		MSept-M-LOct	
END			LOct - Nov		LSept-M-LOct	

REMARKS

*OBS - observed


Muddy Creek
N
0
2
kilometres
River
Eckstall
Rae Lake
Mill Lake
Creek
Muddy
48
49
50
51
52
53
56
57
58
64
65
66
67
68
69
70
71
300
500
1000
1100
1500
1600
2000
3000
3500

NAME OF STREAM MUDDY CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows E. into Ecstall River, Rge 4, Coast Dist.

POSITION 53 129 NW.

LENGTH ___ km WIDTH ___ m DRAINAGE ___ km²

COMPOSITION: BEDROCK ___ BOULDER ___ COARSE ___ FINE ___

SILT & SAND ___ UNCLASSIFIED ___

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ___ m² SPAWNING AREA ___ m²

DISCHARGE (m³/s) ___

TEMPERATURE (°C) ___

BARRIERS OR POINTS OF DIFFICULT ASCENT Impassable cascading rock falls approx. 3.2km from mouth.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	- below falls on small gravel bar.
PINK (EVEN YEAR)	- below falls on small gravel bar.
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ___

GENERAL REMARKS 1947 First time pinks were observed.

1949 - Only limited spawning in this stream.

1953 - Large rainbow and cutthroat taken from lake--popular sportsfishing ground.

1968 - Suggest that stream be included in Ecstall River escapement figures.

1971 - Suggest that stream be discontinued because spawning area is limited and no fish seen.

- Tidally influenced within 90m of falls.
- Between falls and Ecstall River stream is slow running and muddy.
- Mountainous surrounding country.
- Seasonal fluctuations in water levels.
- Predators: bears,wolves and wildfowl.



Sketch of Muddy Creek,

o - Pink Spawning Grounds
~ - Limit of Tidal Influence
∴ - Sandy, Muddy Bottom
// - Impassable Rock Falls
Ecstall River
Lake
Muddy Creek
N
0
1.33
kilometres

ESCAPEMENT RECORD FOR MUDDY CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947					1500	
48						
49						
50					3500	
51					N/0	
52						
53					400	
54						
55						
56						
57						
58						
59					N/0	
60					N/0	
61					N/0	
62					N/0	
63					400	
64					N/0	
65					N/I	
66					N/0	
67					N/0	
68					N/0	
69					200	
70					N/0	
71					N/0	
72					N/0	
73					N/0	
74					N/0	
75					N/0	
76					N/0	
77					N/0	
78					N/0	
79					N/0	
80					N/0	
81					N/I	
82						
83						
84						
85						

TIMING						
ARRIVE						
START					MAug-ESept	
PEAK					M Sept	
END					M-LSept-Oct	

REMARKS N/I - Not inspected


Murder Creek
N
0
2
kilometres
WM
63
62
61
60
59
58
57
56
55
54
53
Murder
Creek
RIVER
Sammon Lake
KISPIOX R
2400
2900
2800
2500
2300
2200
2100
2000
1700
1600
1500
1300
1200
1100
1000
3023
3022
3021
3020
3020F
3019
3018
3017
3017F
3016
3016F
3015F
3014
3013
3012
3011
3010
3007
3006
3005
3005F
3486
3484A
2279
2278
2277
2276
2275
2274
2273
2272
2271
2270
2269
881
878
877
588F
586
584
Bk. A
Bk. B

NAME OF STREAM MURDER CREEK (Beirness Creek,Beirnes Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S. into Kispiox River, North of McCully Creek,Cassiar Dist.

POSITION 55 127 NW.

LENGTH 3.2 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Forestry culvert under Kispiox Road impassable at low flows. Remedial work done in 1965.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- from 0.8km to 3.2km upstream from mouth.
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1952 - Stream not inspected through locals report moderate pink run and light coho run.

1953 - Locals reported light coho and pink runs.

1954 - Locals reported moderate pink run and good coho run.

1955 - " " good pink and moderate coho runs.

1956 - " " very light pink and medium coho runs.

1962 - Minor pink producer on heavy years if adequate water supply.

1965 - Lowered forestry culvert 1.2m to permit easy access to coho but velocities were too greatduring floods to allow salmon passage. Recommended that culvert be replaced by a bridge. Good coho rearing area in deciduous brush growth.

continued.....

continuation

GENERAL REMARKS MURDER CREEK (Beirness Creek,Beirnes Creek)

1965 - Beaver dams removed as required.
1966 - Forestry culvert modified to enable fish to pass by during flood. Also 67.
1969-78 Erosion,silting, scouring and channeling in lower 0.8km.

Seasonal fluctuations in water levels.
Creek runs mostly throughout flat country and is very shallow most of the year.


Coho
Spawning
Area
Kispiox
Valley
Road
Kispiox
River
N
0
2.56
kilometres

Sketch of Murder Creek, 1971.

ESCAPEMENT RECORD FOR MURDER CREEK (Beirnes, Beirness Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			200		75	
48			200		75	
49			400		200	
50			* N/O		N/O *	
51						
52			NOT INSPECTED			
53			"		"	
54			"		"	
55			"		"	
56			"		"	
57			"		N/O	
58			* N/O		* N/O	
59			NOT INSPECTED			
60			N/O		* N/O	
61			25		25	
62			few		N/O	
63			NOT INSPECTED			
64			500			
65			200			
66			200			
67			200			
68			400			
69			25			
70			25			
71			25			
72			25			
73			25			
74			25			
75			25			
76			25			
77			25			
78			200			
79			N/O			
80			200			
81			75			
82						
83						
84						
85						

TIMING						
ARRIVE			Sept			
START			MOct - ENov			
PEAK			E-M Nov			
END			L Nov - Dec			

REMARKS

* Low water prevented fish from entering creek during inspection period


CHISMORE RANGE
ARTHUR PASSAGE
MOUNTAINS
ANGES
Lewis I
Kelp Passage
Henderson Point
Oona River
Oona Pt
Peninsula Pt
BARESIDE MOUNTAIN
Bareside Pt
Rippon Pt
Spiller
COAST LAND DISTRICT RANGE 5
COAST LAND DISTRICT RANGE 4
1671
1461
1459
1669
1667
1679
1681
1683
1685
1687
1689
1691
1693
1695
1697
1699
2422
2423
2424
2430
5938
5937
5936
5935
5934
5933
6786
2206
2205
2204
2203
2202
2201
2200
2199
1751
1753
N
0
2
kilometres

NAME OF STREAM OONA RIVER

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S.E. into Ogden Channel, Rge.5, Coast Dist.

POSITION 53 130 NE.

LENGTH 8 km WIDTH ___ m DRAINAGE ___ km²

COMPOSITION: BEDROCK ___ BOULDER ___ COARSE ___ FINE ___

SILT & SAND ___ UNCLASSIFIED ___

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ___ m² SPAWNING AREA ___ m²

DISCHARGE (m³/s) ___

TEMPERATURE (°C) ___

BARRIERS OR POINTS OF DIFFICULT ASCENT ___

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- Throughout system above tidal influence
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	" "
STEELHEAD	" "

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ___

GENERAL REMARKS 1953 - Brakish water near mouth.
1958 - Light steelhead run noted.
1961 - Early run fish dies unspawned.
1964 - Coho seen but not counted.
1964 - Severe draught, water too low for salmon to migrate upstream very far.
1971 - Stream clearance suggested.
1972 - Evidence of overspawning at peak of run.
1972-80 Series of passable log jams and beaver dams kept open by minor hand work.
1978 - Some pink spawn may have been damaged by heavy September rains.

continued.....

continuation.

GENERAL REMARKS OONA RIVER

1978 - Possible stream bed damage caused by November floods.

1980 - 21000 pinks from upper bridge to swamp.
1000 " above swamp
2000 " in Oscar Creek
some dead pinks found unscarred and unspawned at upper limit of tidal area.Cause of death unknown.

1981 - Oona River is scheduled for a S.E.P.Community Involvement Project to enhance coho.The group monitored stream flows and water temperatures during winter 81/82 and hope to work on the incubation box for coho in the fall of 82.

Good spawning stream --seasonal fluctuations in water levels.
Predators: eagles,seals, bears and wolves.



Sketch of Oona River, 1968.

PORCHER ISLAND
□ - Coho Spawning Grounds
○ - Pink Spawning Grounds
- Trail
- Foot Bridge
- Limit of Tidal Influence
N
0
.40
kilometres
2nd Bridge (Some Boulders)
1st Bridge
Village
Oona River
Odgen Channel

ESCAPEMENT RECORD FOR OONA RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947					7500	
48			750		35000	
49			N/O		15000	
50			N/O		15000	
51					1000	
52					30000	
53					500	
54					15000	
55			400		7500	
56					27000	
57			1500		750	
58			500		2000	
59			OBS		3500	
60			N/I		200	
61			1500		3500	
62			OBS		7500	
63					3500	
64			OBS		750	
65					750	
66			3500		15000	
67			400		750	
68			750		7500	
69			750		3500	
70			N/O		3500	
71			2500		15000	
72			600		13000	
73			250		2200	
74			300		7000	
75			300		4000	
76			300		50000	
77			300		3000	
78			300		10000	
79			250		5000	
80			350		24000	
81			300		4000	
82						
83						
84						
85						

TIMING

ARRIVE			E Sept		MAug-ESept	
START			LSept - EOct		LAug-E Sept	
PEAK			Sept - LOct		M-L Sept	
END			Oct - E Nov		LSept-MOct	

REMARKS N/I - not inspected


N
Price Creek
0
2
kilometres
KITWANGAR INDIAN
RESERVE 1
Kitwanga
585 P
BM 566
Ferry
16
SKEENA
RIVER
NATIONAL
Price Creek
L 3951
L 3950
L 835
L 2392
L 2391
L 834
L 2398
L 3503
L 833
L 836
L 837
L 2891
L 2890
L 3364
L 3949
L 3948
Tum-bah IR 5
Shandilla Lakes
L 1313
L 1318
L 1319
L 1320
L 1317
L 2955
L 1970
L 1316
L 1971
L 832A
L 832
1153±
60
61
WM
WL
RANGES
600
1000
4000
5000
5500
3500

NAME OF STREAM (Price Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows N. into Skeena River approx 1 .6km W.of Kitwanga

POSITION 55 128SE.

LENGTH 32 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- upper reaches
CHUM	- below highway 16 bridge
PINK (ODD YEAR)	- in lower 3.2km
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1947 - No fish seen this year. Reason for absence unknown.

1949 - Slight scouring in lower reaches during most years between 1970-79.

1952 - Public works channelization caused dewatering of the channel and the loss of spawn deposited in that area.Damage rectified because stream reverted to old course.

1955 - Stream changed course for approx 90m near mouth.

1958 - Low water and high temperatures may account for the complete lack of spawners.

1961 - Scouring near culvert. High velocities. Culvert may require baffles to ensure adequate passage of fish 62.

continued.....

continuation.

(Price Creek)

1962 - Scouring at culvert site.
1965 - Highways road bridge construction must have alt ered conditions as 1950 show good pink run and poor "A" runs since .
1966 - Channelling for culvert has made this stream unstable at the mouth and subject to flash floods 67,68,69.
1967 - Considerable scouring and erosion. 68,69,74,78.
1970 - Heavy scouring during high water 71,72,73,75.
1979 - Difficult to impassable because steep-sloped gravel desposit accumulated following floods in Oct. 1978. Suggest that crews reduce slope at outlet and confine water to one channel. 80.

Road culvert construction in 1960. Stream bed was disturbed by Public Works program channellization to stabilize flow. Steep, bouldery gradient. Seasonal low flow in Pink run period. Evidence of erosion at culvert,footing cement eroded.

Seasonal fluctuations in water levels.


Skeena River
Price Creek
□ - Coho Spawning Grounds
○ - Pink Spawning Grounds
▲ - Chum Spawning Grounds
- Road (Highway 16)
- Road Bridge
N
0
5.12
kilometres

Sketch of Price Creek, 1971.

ESCAPEMENT RECORD FOR (Price Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947					N/O	
48					1500	
49					600	
50					3000	
51					2000	
52					6000	
53					2000	
54					3500	
55					1500	
56					750	
57					1500	
58					N/O	
59			OBS		N/O	
60					N/O	
61			6		300	
62					25	
63					N/O	
64					75	
65					55	
66			25	25	25	
67					25	
68			75	N/O	75	
69			25		N/O	
70			25		N/O	
71			25			
72			25		200	
73			25		25	
74			25		25	
75			25		75	
76			25		25	
77			25		25	
78			25		25	
79			N/O		N/O	
80			N/O		N/O	
81			N/O		N/O	
82						
83						
84						
85						

TIMING						
ARRIVE			LSept - Oct		Aug	
START			M Oct	E Sept	EAug-E Sept	
PEAK			L Oct -M Nov	M Sept	L Aug	
END			Nov - Dec	L Sept	Sept - MOct	

REMARKS


SKEENA FOREST
LAKELSE LAKE
Lakelse Lake
Furlong Bay
Catt Pt
Gainey Pt
Squirrel Pt
Mailbox Point
Radio
Lakelse Hot Springs
LAKELSE LAKE PROV PARK
Campsite
Hatchery Creek
Herman Creek
Ena Ck
Ena Lake
End Lake
Andalas Ck
Clearwater Creek
Schulbuckhand Creek
FORCEMAN RIDGE
Onion Lake
Clearwater Lakes
Lakelse IR 25
Lakgass IR 87
TREE FARM LICENCE 41
24
25
26
27
28
29
30
31
32
2660 T L 8282
2662 T L 8283
2674 T L 8276
2676 T L 11370
2677
2678 T L 11128
2679 T L 11127
2680 T L 11126
2665 T L 11146
2664 T L 11144
2663 T L 11145
2672 T L 11135
2684 T L 8278
TL 7724
TL 7725
500
700
1000
1500
N
0
2
kilometres

Schulbuckhand Creek

For more detail see appendix p. 421

NAME OF STREAM SCHULBUCKHAND CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows N.W. into S. end of Lakelse Lake. Rge.5. Coast Dist.

POSITION 54 128 SW.

LENGTH ______ km WIDTH ______ m DRAINAGE ______ km²

COMPOSITION: BEDROCK ______ BOULDER ______ COARSE ______ FINE ______

SILT & SAND ______ UNCLASSIFIED ______

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ m² SPAWNING AREA ______ m²

DISCHARGE (m³/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT ______

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- scattered to 3.2km upstream of lake
CHINOOK	
COHO	- " " "
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ______

GENERAL REMARKS 1951 - Some scouring during flash fall floods.Stream overflows its banks in first few km from mouth lowering water levels and restricting spawning. Cribbing was erected to assist passage of sockeye.

1952 - A few coho migrated upstream, but most were prevented from entering and eggs from 40 fish were transplanted to other streams.

1953 - Breakthrough repaired.

1956 - Large beaver dam removed, also reported 57.

1960 - Considered as a possible site for installation of a hatchery.

1961 - Reports of algae clogging pumps during January incubation period.

continued.....

continuation.

GENERAL REMARKS- SCHULBUCKHAND CREEK

1962 - Most coho stripped for hatchery operation. Cleared considerable debris from stream.
1964 - Estimates based on hatchery fence count.
- Most of run allowed to spawn naturally.
- Logging operations on side hills in upper reaches.
1965 - Past hatchery requirements for sockeye and releases to natural full escapement accounts for comparison fluctuations.
1966 - Hill scarification and clearing for Kemano-Prince Rupert transmission line causing considerable siltation.Problem being attended to because stream is important water source for hatchery.
1967 - Stream re-channelled in September to conserve flow in upper areas.
1974 - Moderate silting, scouring and erosion resulted from October flooding.
1977 - Some channelling near mouth and passable beaver dams.
1980 - Beaver dams 2.4km from mouth rendered stream impassable.



Sketch of Schulbuckhand Creek, 1971.

N
0
.50
kilometres
Schulbuckhand Creek
X - Sockeye Spawning Grounds
☐ - Coho Spawning Grounds
- Marshy Area
- Road (Highway 25)
Canal
Lakelse Lake

ESCAPEMENT RECORD FOR SCHULBUCKHAND CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947	400		75			
48	1500		75			
49	1300		200			
50	600		150			
51	1200		100			
52	1100					
53	750		100			
54	750		75			
55	200		75			
56	75		75			
57	200		75			
58	750		200			
59	750		N/I			
60	1000		25			
61	400		75			
62	400		450			
63	400		N/I		75	
64	3000		75		3	
65	3867		25			
66	3500		63			
67	6800					
68	3500		400			
69	1500		200			
70	200		200			
71	750		200			
72	750		200	200		
73	400		200	N/O		
74	700		200			
75	400		75			
76	750		200			
77	750		75			
78	3500		75			
79						
80	2060		250			
81	2800		100			
82						
83						
84						
85						

TIMING

ARRIVE	LJuly - Aug		Oct	Aug	UNK	
START	E-L Aug		M Oct	E Sept	UNK	
PEAK	E-M Sept		LOct-MNov	M Sept	UNK	
END	MSept - Oct		Dec	Oct	UNK	

REMARKS N/I - not inspected


N
Scotia River
0
10
kilometres
TREE FARM LICENCE 1
BLOCK 1
Kasiks
River
KASIKS MTN
Arden Lake
Arden
Khyex
Lachmach
S K E
R A N
F O R
Telegraph Point
Kwinitsa
Salvus
Aberdeen
Windsor Pt
Ayton
Tyee
BALMORAL
Ecstall
River
Khtada Lake
Khtada
TREE FARM LICENCE
BLOCK 1
MT HAYWARD
1262
Hayward
Cuthbert
Big Falls
Lake
1233
1156
750
1500

NAME OF STREAM SCOTIA RIVER

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows N.W. into Skeena River opposite Telegraph Point, Rge.5, Coast Dist.

POSITION 54 129 SW.

LENGTH 24 km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2

SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	- scattered up to 19km from mouth
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1949 - Good pink and coho runs observed by Columbia Cellulose workers, through no official inspection was made.

1954 - Locals reported fair pink run.

1960 - Stream in a timber sale area and logging expected in 1961.

1961 - Inspection cancelled because of floods, also in 1964.

1961 - Siltation noted near mouth 65,66,68,69.

1966 - Fluctuations in numbers of spawning population depends on extent and timing of inspections.

1970 - Turbid water impaired inspection.



Sketch of Scotia River, 1971.

Telegraph Point
Skeena River
Scotia River
- Coho Spawning Grounds
- Pink Spawning Grounds
N
0
5.12
kilometres

ESCAPEMENT RECORD FOR SCOTIA RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49					10000	
50					OBS	
51						
52						
53						
54					N/I	
55			N/I		7500	
56					OBS	
57					3500	
58					OBS	
59					3500	
60					N/I	
61					N/I	
62					25	
63					N/O	
64					N/O	
65					100000	
66					3500	
67					15000	
68					3500	
69					7500	
70					3500	
71			400		3500	
72			400		3500	
73					3500	
74					750	
75					3500	
76					200	
77					1500	
78					550	
79						
80					500	
81					5000	
82						
83						
84						
85						

TIMING

ARRIVE			Oct		Aug	
START			M Oct		EAug-ESept	
PEAK			M Nov		LAug-MSept	
END			Dec		M-L Sept	

REMARKS N/I - not inspected


Shames River
N
0
2
kilometres
Shames
River
S K E E N
5200
5000
3500
3000
2000
1000
1500
2500
3500
4000
5500
4500
5589
5190
5599
2500
5613
5612
5617
5618
5619
5623
5622
5621
539

NAME OF STREAM SHAMES RIVER

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S.E. into Skeena River, E. of Shames, Rge.5, Coast Dist.

POSITION 55 128 SW

LENGTH ______ km WIDTH ______ m DRAINAGE ______ km²

COMPOSITION: BEDROCK ______ BOULDER ______ COARSE ______ FINE ______

SILT & SAND ______ UNCLASSIFIED ______

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ m² SPAWNING AREA ______ m²

DISCHARGE (m³/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT ______

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	- from mouth of stream upstream for 0.8km
PINK (ODD YEAR)	- " "
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ______

GENERAL REMARKS 1964 - Swift stream subject to scouring between road and confluence with Skeena River

- Too unstable for spawning.
- Few chum carcasses noted off stream mouth in Skeena River.
- Flash flood common. Road and railway bridge sections rip-rapped to prevent washouts.

Seasonal fluctuations in water levels.


SKEENA FOR
Shames River
5164
5190
3500
2000
3000
4000
1000
1500
4508
4509
4510
2624
7592
7535
135
33
31
30
29
28
27
26
25
24
23
5619
5618
5617
5616
5615
5613
5612
5611
Shames
3997
Yellowhead Highway
500
Slough
16
Shames
3657
CANADIAN NATIONAL
Graveyard Pt
T L 9502
SKEENA
TREE FARM LICENCE 1
5499
KIT
WL
Whitebottom Creek
200
03
04
05
06
07
08
09
10
N
Shames Slough
0
2
kilometres

NAME OF STREAM SHAMES SLOUGH

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Skeena River side channel at Shames Rge.5,Coast Dist.

POSITION 54 128 SW

LENGTH km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	- in lower 0.8km of slough
PINK (ODD YEAR)	- " " "
PINK (EVEN YEAR)	- " " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1961 - Good spawning year. Redd sampling proposed for research purposes.

1963 - Heavy siltation resulted in 50% survival of eyed eggs and 10% alevin survival to emergence.

1964 - Passable log jam 0.8km from mouth.

- winter survival questionable due to low flows and subsequent exposure to frost.

Siltation noted most years between 1965 and 81.

1965 - Low winter flows continued to cause desiccation.

continued.....

continuation

GENERAL REMARKS SHAMES SLOUGH

1966 - Spawning area dependent upon annual Skeena River flow.
1974 - Heavy scouring destroyed most of spawn in slough.
Alevin survival ranges between 2% and 4%.
Predators: bears and eagles.

ESCAPEMENT RECORD FOR (Shames Slough)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61	NEW	FILE			87500	
62						
63				75	5500	
64				200	8000	
65				N/O	N/O	
66				OBS	716	
67				75	1500	
68				25	1500	
69				25	750	
70				200	1500	
71				200	1500	
72				400	3500	
73				400	400	
74				400	25	
75				25	400	
76				25	25	
77				75	200	
78				50	25	
79						
80				100	6	
81				100	350	
82						
83						
84						
85						

TIMING						
ARRIVE				Aug	Aug	
START				MAug-ESept	MAug-ESept	
PEAK				M Sept	M Sept	
END				Oct	Oct	

REMARKS


Hardscrabble
Creek
Sand
Shannon
Creek
Skeena River
CANADIAN NATIONAL
CHIMDIMA
INDIA
RESERVE
16
S K
5000
4500
4000
3500
3000
2500
2400
2000
1900
1800
1500
1000
500
3900
4300
3200
L 921
L 5979
L 5980
L 4120
L 4121
L 4780
L 194
L 466
L 922
L 6638
L 1436
L 4779
L 5978
L 4119
L 7932
L 1437
L 72
L 152
N
Shannon Creek
0
2
kilometres

NAME OF STREAM SHANNON CREEK (Nicholson Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S.E. into Skeena River, N.E.of Usk, Rge. 5, Coast Dist.

POSITION 54 128 NE

LENGTH ______ km WIDTH ______ m DRAINAGE ______ km²

COMPOSITION: BEDROCK ______ BOULDER ______ COARSE ______ FINE ______

SILT & SAND ______ UNCLASSIFIED ______

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ m² SPAWNING AREA ______ m²

DISCHARGE (m³/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT ______

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ______

GENERAL REMARKS 1948 - Although not formally inspected, locals report good seedings of pink and coho salmon.

1957 - Inspection too late to enumerate pink spawners.

1960 - Very minor stream

1964 - Considerable gravel scour observed.

1968 - Subject to scouring during flash floods 69,70.

1969 - No salmon observed since 1967 . File closed.

Annual gravel removal required at C.N.R.Bridge.

Bouldery at mouth--steep gradient.

Subject to drying during seasonal drought periods.

ESCAPEMENT RECORD FOR SHANNON CREEK (Nicholson Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			75		200	
48			N/I		N/I	
49						
50						
51						
52						
53						
54						
55			N/I		N/I	
56			N/I		N/I	
57					OBS	
58			N/I		N/I	
59			N/I		N/I	
60			N/0		N/0	
61			N/0		N/0	
62			N/0		N/0	
63			N/I		N/I	
64			N/0		N/0	
65			N/0		N/0	
66			N/0		N/0	
67			N/R		N/R	
68			N/0		N/0	
69			N/0		N/0	
70			FILE CLOSED			
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81						
82						
83						
84						
85						

TIMING						
ARRIVE			UNK		UNK	
START			UNK		UNK	
PEAK			UNK		UNK	
END			UNK		UNK	

REMARKS N/I - not inspected


RANGES
MOUNTAINS
PENINSULA
Naden Islets
Oberg Cove
Beatty Point
Woodworth Lake
Shawatlan Lake
Shawatlan River
Shawatlan Indian Reserve 4
L 397
L 735
Mahlon Lake
Bannock Lake
Butze Rapids
Butze Point
Denise
444
N
Shawatlan River
0
2
kilometres

NAME OF STREAM SHAWATLAN CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows SW into Shawatlan Lake, Rge 5, Coast Dist.

POSITION 54 130 SE

LENGTH 4.8 km WIDTH ______ m DRAINAGE ______ km^2

COMPOSITION: BEDROCK ______ BOULDER ______ COARSE ______ FINE ______

SILT & SAND ______ UNCLASSIFIED ______

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ m^2 SPAWNING AREA ______ m^2

DISCHARGE (m^3/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT Falls above power house approx 4.8km from stream mouth.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- from mouth to power house (see map for specific location)
CHINOOK	- " " "
COHO	- " " "
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM no spawning beds above falls.

GENERAL REMARKS 1948 some bank erosion noted.

- trout plentiful 1949,50

1950 - Some scouring near mouth.

1952 - Scouring noted near power house.

1953 - Discharge from power supply pipeline enters this stream. No habitat deterioration observed.

Good gravel below power house in lower reaches.
Subject to seasonal fluctuation in water levels.
Predators: eagles, trout, seals and bears.

continued.....

continuation.

GENERAL REMARKS SHAWATLAN CREEK

1962 - Coho run late and water too high to estimate numbers, 63,64.
1967 - Abnormal powerhouse shutdown caused low flows and Sept. 23 floods caused erosion,scouring and silting below power house. B.C.Hydro has assured adequate flows to correct problem.
1968 - 5% silting from mouth to powerhouse.
- Stream stabilized by City of Prince Rupert.
1971-77,80 Slight siltation observed.
1981 - The city of Prince Rupert and B.C.Hydro plan to replace dam on Woodworth Lake in 1982. Old dam was deemed unsafe in early 1981. The water was released over a period of 2 days to relieve pressure. Abnormally high flows resulted. Some spawn was disturbed and lost.


Sketch of Shawatlan Creek, 1968.

N
0
.56
kilometres
- Sockeye Spawning Grounds
- Chinook Spawning Grounds
- Coho Spawning Grounds
- Pink Spawning Grounds (When Present)
- Chum Spawning Grounds (When Present)
- Impassable Rock Falls
- Powerhouse
Shawatlan River
Shawatlan Lake

ESCAPEMENT RECORD FOR SHAWATLAN RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947	7500					
48	7500				200	
49	3000				500	
50	4500					
51	5000				2000	
52	5000				500	
53	3500		75	200	400	
54	3500		N/O		200	
55	1500				N/O	
56	3500				N/O	
57	1500				400	
58	1200		25		N/O	
59	3500		N/I			
60	3500		OBS	N/O	N/O	
61	3500		OBS	OBS	OBS	
62	3500		OBS			
63	3500		OBS			
64	3500		OBS			
65	3500		750	N/O	N/O	
66	2000	8	3500			
67	3500	25	750			
68	1500	200	750		400	
69	1500	N/O	750		75	
70	1500	N/O	1500		N/O	
71	2000		250		N/O	
72	1000		500		N/O	
73	800		200			
74	600		300			
75	1000		200			
76	700		200			
77	200		300			
78	400		N/O			
79	250		150			
80	1400		UNK			
81	800		200			
82						
83						
84						
85						

TIMING

ARRIVE	E Sept		Sept - EOct			
START	M Sept	LAug-ESept	Oct	Aug	July-ESept	
PEAK	L Sept	M Sept	L Oct	Sept	LAug-MSept	
END	LSept-Oct	LSept-EOct	Nov	Oct	LSept-Oct	

REMARKS


Shegunia River
N
0
2
kilometres
PROVINCIAL FOREST
BABINE
SKEENA
Shegunia
Collins Lake
Robinson Lake
SKEENA RIVER
KISPAIOX INDIAN RESERVE 1
AGWEDIN INDIAN RESERVE 3
Quan-skum-kaah-mich-mich Indian Reserve 4
SIK-E-DAKH
Kispiox
WM

NAME OF STREAM SHEGUNIA CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows E. into Skeena River at junction of Skeena nd Kispiox Rivers, Cassiar Dist. POSITION 55 127 SW.

LENGTH ____ km WIDTH ____ m DRAINAGE ____ km²

COMPOSITION: BEDROCK ____ BOULDER ____ COARSE ____ FINE ____

SILT & SAND ____ UNCLASSIFIED ____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ____ m² SPAWNING AREA ____ m²

DISCHARGE (m³/s) ____

TEMPERATURE (°C) ____

BARRIERS OR POINTS OF DIFFICULT ASCENT Canyon cascades 3.2km from stream mouth. Passable at optimum water levels; difficult at high and low flows until 1972 when debris washed out.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- near cascades
COHO	- near cascades and sometimes above.
CHUM	
PINK (ODD YEAR)	- lower reaches
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM Good spawning grounds above cascades.

GENERAL REMARKS 1961 - Poor observation conditions.

1965 - Suggested engineering survey of effect of falls on ascending salmon during high velocity flows through gorge.

1966 - Good chinook, coho and steelhead fishing below canyon to mouth.

- Flood caused erosion, fast flows, turbid water and poor observation condit.

1968 - Some silting and scouring resulted from autumn floods, also 78,79,80.

1972 - High water washed out debris in old channel which was causing the river to divert and creating falls. Blasting might keep river permanently in old channel and prevent drying during low flows.

continued.....

continuation.

GENERAL REMARKS SHEGUNIA RIVER.

1973 - River was rechannelled around falls and rerouted to its original stream bed.
Fish may have passed through without notice.
1978 - Cascade should be inspected for possibility of installing a fish ladder to enable salmon to use the spawning grounds above the cascades not presently used.
1981 - Irrigation ditch along side the river monitored regularly because may cause problems for spawning fish.

Generally rough,rocky, streambed with moderately turbulent flows.
Seasonal fluctuations in water levels.
Predators: bear, birds and eagles.


Kispiox River
Skeena River
Shegunia River
Bulkley River
□ - Coho Spawning Grounds
N
0
8.0
kilometres

Sketch of Shegunia Creek, 1975.

ESCAPEMENT RECORD FOR SHEGUNIA RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48			400		200	
49		25	75		750	
50		75	200		400	
51						
52		N/I	N/I		N/I	
53		N/I	N/I		N/I	
54		OBS	N/O		OBS	
55		OBS				
56		OBS			OBS	
57		N/I	N/I		N/I	
58		N/I	N/I		N/I	
59		N/I	N/I		N/I	
60		N/O	N/O		N/O	
61		N/O	N/O		N/O	
62		N/O	N/O		N/O	
63		N/O	N/O		N/O	
64		N/O	N/O		N/O	
65		400				
66		25	400			
67		75				
68		25	400			
69			300			
70			300			
71		50	200			
72		N/R	N/R			
73			N/O			
74			N/O			
75						
76		N/O	N/O			
77		N/R	N/R			
78			OBS		75	
79			OBS		25	
80		25	150		350	
81		25	75		400	OBS
82						
83						
84						
85						

TIMING

ARRIVE		July-Aug	Sept		Aug	
START		E Aug	L Sept-Oct		M Aug	
PEAK		L Aug	EOct-MNov		LAug-ESept	
END		E Sept	E-L Nov		Sept	

REMARKS N/I - not inspected


Silver Creek
0
2
kilometres
N
Tuck Inlet
Tuck Narrows
Naden Islets
Beatty Point
Scissors
MOUNTAINS
RANGE

NAME OF STREAM SILVER CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S.W. into Prince Rupert Harbour at Tuck Narrows, Rge.5, Coast Dist.

POSITION 54 130 SE.

LENGTH ______ km WIDTH ______ m DRAINAGE ______ km²

COMPOSITION: BEDROCK ______ BOULDER ______ COARSE ______ FINE ______

SILT & SAND ______ UNCLASSIFIED ______

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ m² SPAWNING AREA ______ m²

DISCHARGE (m³/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT Impassable rock falls 4.8km upstream from mouth.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- from 0.8 to 4.0km upstream from mouth
CHUM	- lower reaches
PINK (ODD YEAR)	- from 0.8 to 4.0km upstream from mouth
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ______

GENERAL REMARKS 1948 - Stream changed course for approx 274m near mouth.

Some erosion evident in upper portions of this stream following heavy rains.

Stream bed rocky beyond spawning limits. Sufficient gravel to accomodate run.

Seasonal fluctuations in flows.

Predators: bears, seals, wolves and eagles.



Sketch of Silver Creek, 1971.

N
0
.84
kilometres
▯ - Coho Spawning Grounds
○ - Pink Spawning Grounds
▲ - Chum Spawning Grounds
✕ - Passable Log Jam
// - Impassable (Rock) Falls
Silver Creek
Tuck Inlet
Tuck Narrows
To Prince Rupert Harbour

* Logged about 40 yrs ago along both banks

ESCAPEMENT RECORD FOR SILVER CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947				3500	3500	
48				3500	7500	
49				1500	7000	
50				6000	4000	
51				1000	3000	
52				300	1500	
53				1000	1000	
54				2000	5000	
55				25	25	
56			10	6	35	
57			200	N/O	400	
58				100	50	
59				25	25	
60				75	200	
61				400	1500	
62				25	200	
63				200	1500	
64				200	1500	
65				NOT INSPECTED		
66			1500	75	3500	
67				25	200	
68			N/O	N/O	3500	
69			400	N/O	750	
70			N/O	N/O	400	
71			UNK	N/O	1200	
72			UNK	N/O	500	
73			UNK	N/O	200	
74			UNK		350	
75			UNK		300	
76			UNK		250	
77			UNK		350	
78			UNK	50	1000	
79			UNK		125	
80			UNK		800	
81					1000	
82						
83						
84						
85						

TIMING

ARRIVE			Oct	Aug	Aug	
START			Oct	Sept	Sept	
PEAK			Nov	Sept	MSept-MOct	
END			Nov	Oct	LSept-Oct	

REMARKS


SKEENA
Usk
Ferry
CANADIAN NATIONAL
16
KITSELAS
INDIAN RESERVE
KLEANZA CREEK PROV PARK
KLEANZA CREEK PROVINCIAL PARK
Sawmill
Tunnels
Tunnel
Singlehurst Creek
Yellowhead Highway
Kleanza Creek
Noble Creek
Noble Five Lake
Kitselas
KLEANZA MOUNTAIN
N
Singlehurst Creek
0
2
kilometres

NAME OF STREAM SINGLEHURST CREEK (Swede Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S.W. into Skeena River, N. of Kleanza Creek, Rge.5, Coast Dist.,

POSITION 54 128 NE

LENGTH ______ km WIDTH ______ m DRAINAGE ______ km²

COMPOSITION: BEDROCK ______ BOULDER ______ COARSE ______ FINE ______

SILT & SAND ______ UNCLASSIFIED ______

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ m² SPAWNING AREA ______ m²

DISCHARGE (m³/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT ______

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- upper reaches
CHUM	- near mouth
PINK (ODD YEAR)	- lower 0.8km
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ______

GENERAL REMARKS 1951 - Low water prevented fish from entering stream.

1956 - Low water conditions in 58,62.

1961 - Silting and scouring.

- Numerous coho fry sighted also 65,66.

- Beaver activity noted 62,78,80,81 cleared as required.

1965 - Culvert installed to accomodate relocated section of Highway 16.

1966 - Suggest closing file because stream virtually unused.

1969 - File closed

1978,80 Successful fisnway built to bypass culvert. Formerly an obstacle when highway 16 rerouted.

Subject to seasonal flow rates. Predators: bears and eagles.

ESCAPEMENT RECORD FOR SINGLEHURST CREEK (Swede Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48				25	1500	
49				N/O	N/O	
50				50	600	
51				N/O	N/O	
52					500	
53						
54					200	
55					N/O	
56					N/O	
57					N/O	
58					N/O	
59					N/I	
60					N/O	
61					N/O	
62					N/O	
63					N/O	
64					N/O	
65					N/O	
66					N/O	
67					N/O	
68					N/O	
69					N/O	
70						
71						
72						
73						
74						
75						
76						
77						
78			75		400	
79						
80			55		4	
81			40		25	
82						
83						
84						
85						

TIMING						
ARRIVE			Sept		Aug	
START			L Sept	E Sept	M Aug	
PEAK			M Oct	M Sept	L Aug	
END			Nov	L Sept	Sept	

REMARKS


Skeena River
Main Stem
0
20
kilometres
N
Terrace
Lean-To Mtn
Kitsumkalum
Lakelse
Mt Cell
Kitimat
Kitimat River
Exstew River
Kitnayakwa River
Exchamsiks River
Kasiks River
Khyex River
Ecstall River
Scotia River
Williams Creek
Zymacord
Kleanza Creek
Kitimat Mtn

NAME OF STREAM SKEENA RIVER (Mainstem)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S.W. into Chatham Sound, Rge. 5, Coast Dist.

POSITION 54 130 SE

LENGTH km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- Selma Island, Terrace
COHO	
CHUM	- Remo Bar, Hells Gate Slough
PINK (ODD YEAR)	- Selma Island, Terrace, Extew River.
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1957 - Too late for inspection.
1965 - Development of river for log towing investigated.
Dredging proposed close to main stem spawning area.
1966 - Normal siltation in estuary and side channels also in 68,69,77.
1967 - Shifting gravel in Split Mountain area due to high velocities.
1972 - Reports indicate 1,000,000 pinks in main stem,but 50% of spawn lost
when river level dropped. Effective spawning recorded as 500,000
1981 - Silting and erosion.
Seasonal fluctuations in water levels.
Predators: seals, eagles and bears.

ESCAPEMENT RECORD FOR SKEENA RIVER (Main Stem)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65				3500	125000	
66				2000	25000	
67					90000	
68					150000	
69					35000	
70					20000	
71					50000	
72				1500	500000	
73				1500	25000	
74				1500	10000	
75		1500		1500	200000	
76		1500		750	7500	
77		3500		750	35000	
78						
79		200		1000	20000	
80						
81		500		200	20000	
82						
83						
84						
85						

TIMING

ARRIVE		July		July - Aug	E Aug	
START		M Aug		E Aug	M Aug	
PEAK		L Aug		M-L Aug	L Aug	
END		Sept		E-M Sept	Sept	

REMARKS


Sockeye Creek
N
0
2
kilometres
Kulspai IR6
TL 9508
Terrace Airfield
Thunderbird
Sockeye Creek
Williams Ck
Muller Bay
LAKELSE LAKE PROVINCIAL PARK
LAKELSE LAKE
Furlong Bay
Furlong Creek
Mount Herman
Herman Lake
Hai Lake
Campsite
Gainey Pt
LAKELSE LAKE
SKEENA FOREST
CN
25
2 Lines
Gate
Picnic

NAME OF STREAM SOCKEYE CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S. into Williams Creek, N. of Lakelse Lake, Rge.5, Coast Dist. POSITION 54 128 SW.

LENGTH 8 km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- scattered for 8.0km upstream from junction with Williams Cr.
CHINOOK	
COHO	- scattered for approx.11.0km upstream from junction with Williams Creek.
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1959 - A minor tributary of Williams Creek used only intermittently. Estimates usually included in Williams Creek files,63.

1961 - Williams Creek hatchery program geared to handle run as required.

1962 - Lower 0.8km of stream below Park road culvert has slow flow and heavy mud deposits.66.

1965 - Logging by Pohle Co in upper watershed has denuded that section of stream. Ranger ordered cleanup by end of May 66 otherwise prosecution pending.

continued.

continuation

GENERAL REMARKS COHO CREEK

1966 - Adjacent low lands flooded in October.
- Channel improvement conducted in upper reaches to improve flow and make area more accessible to coho spawners.

1968 - Silting noted 69,74.
1974 - Scouring and erosion noted following fall floods

Log debris scattered throughout, and removed as required 1966-78
Seasonal fluctuations in water levels.
Predators: bears.



Sketch of Sockeye Creek, 1971.

Sockeye Creek
Williams Creek
▢ - Coho Spawning Grounds
✕ - Sockeye Spawning Grounds
△ - Chinook Spawning Grounds
✕ - Road bridge
╱ - Road
N
0
5.12
kilometres

ESCAPEMENT RECORD FOR SOCKEYE CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947	400		400			
48	750		75			
49	500		200			
50	200		200			
51	500		200			
52	N/I		N/I			
53						
54	N/I		N/I			
55	N/I		N/I			
56	N/I		N/I			
57	N/I		N/I			
58	N/I		N/I			
59	*		*			
60						
61	*		*			
62	25					
63	*		*			
64						
65	*		1500			
66	2000	25	400			
67	75					
68	400	25	400			
69	1500	25	400			
70	1500	25	400			
71	750	25	400			
72	400	N/O	400			
73	400	N/O	200			
74	200	N/O	200			
75	400		200			
76	750		200			
77	75	N/O	75			
78	400		200			
79						
80						
81	500		175			
82						
83						
84						
85						

TIMING

ARRIVE	Aug	Aug	Oct			
START	MAug-ESept	Aug - E Sept	L Oct			
PEAK	E-M Sept	M Sept	M Nov			
END	L Sept-Oct	Sept - Oct	Dec			

REMARKS

* Estimates included with Williams Creek

N/I - not inspected


84
85
86
87
88
89
90
91
97
96
95
94
93
92
91
90
89
88
87
86
FOREST
ALASTAIR LAKE
Alastair Peak
Icefield
Southend Creek
TREE FARM LICENCE
2700
2100
5200
5800
4200
2800
700
1400
1900
2000
4000
4700
5600
5000
3500
N
Southend Creek
0
2
kilometres

NAME OF STREAM SOUTHEND CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows into Allistair Creek.

POSITION 54 129 SE.

LENGTH 2.4 km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Impassable falls approx. 30m high located about 2.4km upstream from bifurcation on both South and East forks.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- scattered to falls on both forks at at distance of approx 1.6km
CHINOOK	
COHO	- to falls on both forks.
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1948 - Some evidence of late spawners digging up previously spawned eggs.

1954 - Coho seen off stream mouth, also in 56

1957 - Few coho jacks seen

- Beaver dams removed on East fork.

- Heavy losses expected due to mid season low flows.

1960 - Sockeye run consisted of 40% male, 50% female and 10% jacks.

1961 - "Green" sockeye observed in L. Sept, but water conditions prevented adequate follow up observation.

1962 - Sand deposits near mouth also in 65.

continued.....

continuation.

GENERAL REMARKS SOUTHEND CREEK.

1962 - Main producer of Allistair system sockeye runs.
1963 - Inspection made at mid-point of run. Expect final estimates of 9000 to 10000 spawners.
1964 - Predominantly females.
1965 - Numerous small green 4/2 female sockeye observed.
1966 - Measurements scale and atholith data taken by Fisheries Research Board.
1968 - Silting, scouring and erosion noted following periods of high water, 69,70,77.
1972 - Coho estimates based on reports from local sports fishing guides.

Seasonal fluctuations in water levels--clear run-off,mountainous surroundings.
Predators: bears,wolves,seals,eagles.


Allistair Lake
× - Sockeye Spawning Grounds
□ - Coho Spawning Grounds
// - Impassable Falls
Southend Creek
N
0
2.56
kilometres

Sketch of Southend Creek, 1971.

ESCAPEMENT RECORD FOR SOUTHEND CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48	15000					
49	10000					
50	8000					
51	10000					
52	12000					
53	12000					
54	15000					
55	15000					
56	15000					
57	35000					
58	35000					
59	15000					
60	3000					
61	12500					
62	8445					
63	7500					
64	1500					
65	7500					
66	8000		1500			
67	7500		3500			
68	7500		7500			
69	1500		750			
70	750		750			
71	750		750			
72	2500		750			
73	2500		400			
74	1000		400			
75	400		N/I			
76	1500		N/I			
77	3500		N/I			
78	8000		75			
79						
80	10500		N/I			
81	400		N/O			
82						
83						
84						
85						

TIMING						
ARRIVE	Aug		Sept - Oct			
START	MAug - ESept		MSept - MOct			
PEAK	M Sept		E-M Oct			
END	M-L Oct		Dec			

REMARKS


Sparkling Creek
0
10
kilometres
N
Ecstall
River
Herman L.
Big Falls L.
Madeline L.
Kumealon L.
Murray Pt.
Baker Inlet
Griffon Pt.
Alva L.
Kxngeal L.
Kxngeal Inlet
Nordness Pt.
G R E N V I L L E
C H A N N E L
P I T T
K I A T
Lower L.
Ecstall L.
Harrier
Ship Anch.
East Inlet
Foda L.
Studio L.
Gavel L.
Buckland Cr.
Buckland L.
Simpson Rapids
Saunders L.
MT. SAUNDERS
Douglas Pt.
Lowe L.
Rapids Falls
Woerz L.
Quaal
Aley L.
ANCHOR CONE
Hopkins Pt.
Lowe In.
Gamble L.
Belowe L.
SYLVAN
MT. PATTERSON
Bluff
Angel Pt.
Ala Pass
Lagasse Lagoon

NAME OF STREAM SPARKLING CREEK (Clear Water Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows N. into Ecstall River, W. of Johnston L., Rge.4., Coast Dist.

POSITION 53 129 NW.

LENGTH km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Impassable rock fall 2.0km from stream mouth.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	- lower 1.2km of this stream and below falls for 182m.
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1974 - Recommend any spawning escapements be included in Ecstall R.
1977 - High turbidity hampered observation.
1978 - Heavy rains in November may have damaged stream bed.

Muddy bottom for 1.6km from Ecstall River junction. Mostly bouldery, rough gravel. Subject to seasonal fluctuations in water levels. Poor spawning stream.


- Limit of Tidal Influence
- Falls
Ecstall River
Clear Water
Some Gravel
* Most of Bottom Compsed of Rough, Bouldery Substrate
Sparkling Creek
(Clearwater Creek)
N
0
3.33
kilometres

Sketch of Sparkling Creek.

ESCAPEMENT RECORD FOR SPARKLING CREEK (Clear Water Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947				3500	3500	
48				75	25	
49				500	2000	
50					N/O	
51				300	600	
52				2500	5000	
53					200	
54				2	20	
55				25	1500	
56				N/O	N/O	
57					1500	
58				100	100	
59					750	
60				N/O	N/O	
61				N/O	N/O	
62					N/O	
63					750	
64				N/O	N/O	
65				N/I	N/I	
66			N/O	N/O	N/O	
67				N/O	200	
68				N/O	N/O	
69				N/O	N/O	
70				N/O	N/O	
71			UNK		300	
72				N/O	N/O	
73			N/O	N/O	N/O	
74			N/O	N/O	N/O	
75			UNK	N/O	N/O	
76			N/O	N/O	N/O	
77			N/O	N/O	N/O	
78			N/O	N/O	N/O	
79			UNK	N/O	N/O	
80			UNK	N/O	N/O	
81			UNK	N/O	N/O	
82						
83						
84						
85						

TIMING						
ARRIVE						
START				Aug	Aug	
PEAK				Sept	Sept	
END				Oct	Oct	

REMARKS


PASSAGE
ARTHUR
PASSAGE
Francis Point
Chalmers Anchorage
Elliott Island
Hinsam Passage
Lawson Harbour
Break Island
Herbert Reefs
Lewis Island
Kelp Passage
Godfrey Point
RANGE
RANGES
CHISMORE RANGE
Salt Lagoon
Lewis I
Oona River
Oona Pt
N
Spiller River
0
2
kilometres

NAME OF STREAM SPILLER RIVER

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows N.E. into Chismore Passage, Porcher Island, Rge. 5, Coast Dist.

POSITION 54 130 SE.

LENGTH 3.2 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- well dispersed throughout available gravel area.
CHUM	
PINK (ODD YEAR)	- well dispersed throughout available gravel area.
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1950 - Good gravel bed.

1953 - Slough-like, marshy grasslands near mouth.
- Water highly discoloured.
- Tidal influence extends for o.8km upstream from mouth.
- Preponderance of femal pinks observed.

1961 - Early spawners died without spawning dur to severe low flows;however, flows were satisfactory for the bulk of the run to spawn successfully.

1965 - Low water delayed upstream migration.

1967 - Minor silting noted most years between 1967 and 1980.

continued.....

continued

GENERAL REMARKS SPILLER RIVER.

1972 - Some evidence of over spawning.
- Other streams on Porcher Island flowing into Chismore Pass also received good seeding of pink slamon.

1973 - Pink escapement figures include the 150 fish spawning near the mouths of the 3 creeks on Hunts Point.

1976 - Better run than expected.

1978 - Heavy November rains may have caused some stream bed damage resulting in egg loss.

1981 - Coho fry observed. Good rearing conditions in side channels and small tributaries.

Seasonal fluctuations in water levels. Headwaters swampy.
Predators: seals,trout, wolves, eagles and bears.


Chismore Passage
PORCHER ISLAND
Spiller Creek
○ - Pink Spawning Grounds
□ - Coho Spawning Grounds
∿ - Limit of Tidal Influence
Old Bridge
* Fair Gravel for 4.0-4.8 km from Mouth
N
0
.40
kilometres

Sketch of Spiller River.

ESCAPEMENT RECORD FOR SPILLER RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947					7500	
48					7500	
49					15000	
50					10000	
51					1500	
52					40000	
53					500	
54					1500	
55					15000	
56					4000	
57					6000	
58					1500	
59					3500	
60					1500	
61					3500	
62					3500	
63					1500	
64					3500	
65					1500	
66			1500		7500	
67			75		750	
68			75		7500	
69			200		1500	
70			N/0		3500	
71			UNK		7000	
72			UNK		11000	
73			UNK		350	
74			UNK		1000	
75					400	
76					5000	
77					200	
78					3000	
79					200	
80					* 4200	
81			UNK		200	
82						
83						
84						
85						

TIMING						
ARRIVE					M Aug	
START			MSept-EOct		MAug-MSept	
PEAK			LSept-MOct		M-L Sept	
END			M-L Oct		M Sept-Oct	

REMARKS

* Includes 1100 dead unspawned pinks in good condition. Reason for mortalities unexplained. Similar occurrences in Humpback Bay Creek and Oona River.


Kitsumkalum Lake
1400
1426
Treston Lake
1423
1421
1800
1428
1427
1425
Creek
65
64
63
62
61
60
59
58
57
Star Creek
NELSON RIVER
TIMAT
Star Creek
N
0
2
kilometres

NAME OF STREAM STAR CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows E. into Kitsumkalum River, S. of Kitsumkalum Lake, Rge.5, Coast Dist. POSITION 54 128 NW.

LENGTH 6.4 km WIDTH ___ m DRAINAGE ___ km²

COMPOSITION: BEDROCK ___ BOULDER ___ COARSE ___ FINE ___

SILT & SAND ___ UNCLASSIFIED ___

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ___ m² SPAWNING AREA ___ m²

DISCHARGE (m³/s) ___

TEMPERATURE (°C) ___

BARRIERS OR POINTS OF DIFFICULT ASCENT ___

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- scattered for about 4.8km upstream from mouth.
CHUM	
PINK (ODD YEAR)	- lower river when present.
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ___

GENERAL REMARKS Generally only observed to Columbia Cellulose Company Bridge, approx. 0.8km from mouth.

1953 - Cleared canyon blockage.

1954 - High and turbid water prevents accurate observation and assessments in most years.

1956 - Columbia Cellulose Co. cleared log jam at highway bridge.

1961 - Flash flood resulted in considerable silting, scouring and erosion, 69-80. Columbia Cellulose restored channel to contain flow in bridge vicinity.

1964 - erosion noted.

1965 - In C.C.C. tree farm area Kalum -- block mountain valley -- silty runoff-- minor tributory--sandy fine gravel--bouldery -- thick underbrush above bridge. Rocky stream, swift flow. Flash floods and extremely silty water conditions common.

ESCAPEMENT RECORD FOR STAR CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53			200		OBS	
54			N/I			
55			N/O			
56			N/O			
57			OBS			
58			OBS			
59			N/I			
60			N/O			
61			400			
62			N/O			
63			N/O			
64			N/O			
65			N/O			
66		25	75			
67		N/O	N/O			
68			N/O			
69			N/O			
70			N/O			
71			N/O			
72			N/O			
73			N/O			
74						
75						
76			25			
77			25			
78			25			
79			N/I			
80						
81			50		200	
82						
83						
84						
85						

TIMING						
ARRIVE			Oct		Aug	
START		Aug	Nov		E Sept	
PEAK		Aug	M Nov		M Sept	
END		Sept	Dec		L Sept	

REMARKS N/I - not inspected


N
Stephens Creek
0
2
kilometres
KISPIOX
RIVER
Club Lake
Stephens Lake
Stephens Creek
2000
2200
2100
2500
2800
3300
3700
3500
4000
3800
3200
3600
3100
1900
2700
2400
3400
2900
2600
2300
3000
2100
1800
1700
1600

NAME OF STREAM STEPHENS CREEK.

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S.E. into Kispiox River, Cassiar District.

POSITION 55 128 NW.

LENGTH 2.4 km WIDTH ___ m DRAINAGE ___ km²

COMPOSITION: BEDROCK ___ BOULDER ___ COARSE ___ FINE ___

SILT & SAND ___ UNCLASSIFIED ___

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ___ m² SPAWNING AREA ___ m²

DISCHARGE (m³/s) ___

TEMPERATURE (°C) ___

BARRIERS OR POINTS OF DIFFICULT ASCENT ___

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- scattered spawning throughout stream.
CHINOOK	- " " "
COHO	- " " "
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ___

GENERAL REMARKS 1954 - Beaver dam removed 0.8km from Stevens Lake.
1960 - Silting, also in 65.
1961 - Competition for spawning space between chinook and sockeye.
- Sockeye have difficulty ascending Gitwangulf Canyon when flows are low.
1967 - 2500 sockeye and 500 coho through area in L. Sept.

Obstructions removed as required to ensure passage of fish.
Seasonal fluctuations in water levels.
Predators: bears.



Sketch of Stephens Creek.

Stevens
Lake
Kispiox
River
Stevens
Creek
□ - Coho Spawning Grounds
× - Sockeye Spawning Grounds
△ - Chinook Spawning Grounds
N
0
1.0
kilometres

ESCAPEMENT RECORD FOR STEPHENS CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947	579	OBS	OBS			OBS
48	25	N/O	25			
49	10	200	150			
50		100	100			
51	200		400			
52	50	N/O	200			
53		200	50			
54	N/O	200	200			
55	N/O	200	200			
56	25	25	400			
57		75	200			
58		200	400			
59		400	75			
60		2	3500			
61	1500	750	1500			
62	17	N/O	N/O			
63	75		200			
64	75	25	750			
65	75	25	750			
66	400	200	1500			
67						
68		N/O				
69	750		750			
70	400		750			
71	400		750			
72	400		750			
73	400	25	200			
74	400	25	200			
75	200	25	200			
76	200		400			
77	400	400	200			
78						
79	N/O	N/O	N/O			
80	* 100	* 250	* 150			
81	N/I	N/I	N/I			
82						
83						
84						
85						

TIMING						
ARRIVE	L Aug	Aug	Sept			
START	E Sept	M-L Aug	Sept - EOct			
PEAK	M Sept	E-M Sept	MSept-M Oct			
END	LSept-MOct	L Sept	LSept-L Oct			

REMARKS

* Figures obtained from surveys conducted by B.C. Fish & Wildlife.

N/I - not inspected


L 3940
IR 50
Tp 1
Lizzie Hill
WORK CHANNEL
TSIMPSEAN PENINSULA
Trail Bay
Nnaxtoalk Lake
PORT SIMPSON
Grace Point
George Hill
Lamo Point
Bay
Bath Point
Port Simpson
Tsimpsean
Mount Griffin
1364
Mount Ben
1920
Stumaun
Creek
INDIAN RESERVE
RANGES
MOUNTAINS
52
51
50
49
48
47
46
45
44
43
42
41
21
22
23
16
15
14
13
12
11
10
3
4
2
1
500
400
300
200
100
1000
1500
2000
1300
N
Stumaun Creek
0
2
kilometres

NAME OF STREAM STUMAUM CREEK (Simpson Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows into Stumaun Bay, Head of Port Simpson.

POSITION 54 130SE.

LENGTH km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Passable but logging debris scattered throughout. Recommend removal.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	- even throughout entire system
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1978. Water levels extremely low during migration, normal during spawning. Some egg digging in lower end. Good amount of Coho fry seen in this stream. Project with village should be initiated next Spring to remove logging debris.

1979. If funds unavailable to hire people, patrolmen should clean as much logging devris as possible prior to the return of the 1980 pinks.

ESCAPEMENT RECORD FOR STUMAUN CREEK (Simpson Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78					11000	
79					N/O	
80						
81						
82						
83						
84						
85						

TIMING						
ARRIVE					Aug 10	
START					Aug 15	
PEAK					Sept15	
END					Sept 30	

REMARKS N/O-not observed


KHUTZEYMATEEN
INLET
Walskakul Point
McGregor Point
Tsampanaknok Bay
TSAMPANAKNOK CREEK
VL
RANGES
MOUNTAINS)
63
62
61
60
59
58
57
56
55
54
4
25
26
27
28
29
30
31
32
3500
3000
2500
2000
1800
1000
600
500
400
200
100
0
N
Tsampanaknok Creek
0
2
kilometres

NAME OF STREAM (Tsamspanaknok Bay Creek) (Sam Bay)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flowing into Khutzeymateen Inlet.

POSITION 54 130NE

LENGTH km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	- in lower reaches only
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1978 - Water levels normal. Helicopter logging to take place in the area later this fall. 1 chain leave strip to be left for stream bank protection.

Checked stream to 2.5km from mouth -- no obstructions.

ESCAPEMENT RECORD FOR (Tsamspanaknok Bay Creek) (Sam Bay)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78					800	
79						
80						
81						
82						
83						
84						
85						

TIMING						
ARRIVE					Aug.7	
START					Aug. 15	
PEAK					Aug. 30	
END					Sept.15	

REMARKS


Useless Creek
0
2.0
kilometres
N
PORCHER ISLAND
Useless Creek
Useless Bay
Useless Point
Refuge Bay
Table Point
Pearce Pt
Goble Pt
Knox I
Gull I
Barrett I
Morrell Point
Spruce Hill
BELL RANGE
Porcher Creek
Foote Creek
Hunts Inlet
Hunt Inlet

NAME OF STREAM USELESS CREEK (Big Useless Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows N.W. into Useless Bay, Edye Passage, Rge. 5, Coast Dist

POSITION 54 130 SW.

LENGTH 3.2 km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	- from mouth 3.2km
PINK (EVEN YEAR)	- from mouth 3.2km
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1953 - Dark, brackish water during inspection period. Small windfalls 0.8km upstream and above.

1955 - Brakish water noted.

1957 - Small brush jams 0.8km to 3.2km upstream, prevent scouring of spawning gravel and do not restrict fish.

1961 - About 2000 early pinks died unspawned due to insufficient flows. Rain in L. Aug brought the main run in and allowed successfull spawning.

1963 - Low flows hampered upstream migration and caused death of some fish prior to spawning.

continued.....

continued.

GENERAL REMARKS (Big Useless Creek)

1964 - Spawning high on banks and prolonged dry,cold,weather could result in desiccation of these eggs.
1966 - Poor visibility during stream inspection.
1973,1967 Frequent water discolouration and turbidity hamper accurate assessments of salmon stocks.
1976 - Better run than expected.
1977 - Heavy rains in late September.
1978 - Heavy rains in early November caused some stream bed damage.
1979 - Slight siltation noted.

Stream has good gravel sections. 3.2km of suitable gravel in all.
Seasonal fluctuations in water levels and flash floods during freshets.
Predators: hair seals, eagles, wolves.


Edye Passage
Useless Bay
Extent of Tidal Influence
PORCHER ISLAND
o - Pink Salmon Evenly Throughout
- Coho Observed
N
Useless Creek (Big)
0
3.2
kilometres

ESCAPEMENT RECORD FOR USELESS CREEK (Big Useless Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947					3500	
48					3500	
49					7000	
50					4000	
51					1500	
52					30000	
53					400	
54					5000	
55					1500	
56					11000	
57					6000	
58					3000	
59					3500	
60					3500	
61					7500	
62					35000	
63					7500	
64					7500	
65			N/0		200	
66			1500		3500	
67			75		2500	
68			N/0		3500	
69			750		1500	
70			N/0		200	
71			UNK		1800	
72			UNK		7000	
73			200		1425	
74			UNK		2500	
75			UNK		4000	
76			UNK		15000	
77			UNK		3000	
78			UNK		1500	
79			UNK		1500	
80					2000	
81			UNK		850	
82						
83						
84						
85						

TIMING						
ARRIVE					M Aug	
START					MAug-MSept	
PEAK					M-L Sept	
END					LSept-MOct	

REMARKS


PRESCOTT ISLAND
Parry I
Snuff Islet
PRESCOTT PASSAGE
ARTHUR ISLAND
View Pt
Morrell Pt
EDYE PASSAGE
Gorie Pt
Useless Bay
Useless Pt
Edwin Pt
Edye Pass Mine
HENRY ISLAND
Harbour
Little Useless Creek
BELL RANGE
N
Little Useless Creek
0
2
kilometres

NAME OF STREAM USELESS CREEK (Little Useless Creek)

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows N.W. into Useless Bay, Edye Passage, Rge.5, Coast Dist.

POSITION 54 130 SW.

LENGTH ______ km WIDTH ______ m DRAINAGE ______ km^2

COMPOSITION: BEDROCK ______ BOULDER ______ COARSE ______ FINE ______

SILT & SAND ______ UNCLASSIFIED ______

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ m^2 SPAWNING AREA ______ m^2

DISCHARGE (m^3/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT Impassable rock falls on right hand fork 0.8km upstream.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	- evenly dispersed from 0.8km - 1.6km from stream mouth.
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ______

GENERAL REMARKS 1961 - Lack of water delayed upstream migration.
1965 - Lack of water may have affected run size.
1967 - A few pinks use the stream depending upon water levels.
1981 - Low water conditions made creek impassable.
Flows through swampy area, discolored,brakish water.
Subject to seasonal fluctuations in water levels and flash floods.
Predators: eagles, seals and wolves.



Useless Creek (Li ttle)

Edye Passage
Useless Bay
PORCHER ISLAND
Little Useless Creek
No Gravel, only Boulders
● - Pink Spawning Grounds
〃 - Impassable Falls
N
0
2.0
kilometres

ESCAPEMENT RECORD FOR USELESS CREEK (Little Useless Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
					1500	
1947					7500	
48					2000	
49					6000	
50					1000	
51					15000	
52					3500	
53					2500	
54					3500	
55					1500	
56			400		750	
57					1200	
58					750	
59					200	
60					1500	
61					1500	
62					400	
63					25	
64					25	
65					25	
66					N/O	
67					400	
68					750	
69					25	
70					UNK	
71					UNK	
72					N/O	
73					N/O	
74					N/O	
75					N/O	
76					UNK	
77					UNK	
78					N/O	
79					N/O	
80						
81						
82						
83						
84						
85						

TIMING						
ARRIVE			Oct		LAug-ESept	
START			Oct		M Sept	
PEAK			Nov		LSept-EOct	
END						

REMARKS


LAKELSE
RIDGE
ASH
A T
R A N G E
M O U N T A I N S
White Creek
Creek
Mount Johnstone
Icefield
3787
4000
4500
5066
4943
6059 Mount
2655 TL 11134
2654 TL 11123
2653 TL 11120
2659 TL 8285
2658 TL 8284
5976
5973
5972
5971
5974
5240
5239
C
B
A
500
1000
1200
1500
2000
3000
15
16
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
N
White Creek
0
2
kilometres

NAME OF STREAM WHITE CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows N. into Lakelse River, Rge. 5, Coast Dist.

POSITION 54 128SW.

LENGTH ______ km WIDTH ______ m DRAINAGE ______ km²

COMPOSITION: BEDROCK ______ BOULDER ______ COARSE ______ FINE ______

SILT & SAND ______ UNCLASSIFIED ______

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ______ m² SPAWNING AREA ______ m²

DISCHARGE (m^3/s) ______

TEMPERATURE (°C) ______

BARRIERS OR POINTS OF DIFFICULT ASCENT ______

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	- scattered throughout lower section
PINK (EVEN YEAR)	- " " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ______

GENERAL REMARKS 1980/81 Some scouring and silting in lower end. Channel changing below logging road.

ESCAPEMENT RECORD FOR WHITE CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80					30	
81					50	
82						
83						
84						
85						

TIMING

ARRIVE					Aug	
START					Aug 15	
PEAK					Aug 30	
END					Sept	

REMARKS


SKEENA
MOUNTAINS
FOREST
RANGES
BORNITE RANGE
TERRACE
TREE FARM LICENCE 1
BLOCK 3
KLEANZA MTN
COPPER MTN
MT THORNHILL
KITSELAS MTN
LEAN-TO MTN
MT GARLAND
MT LAYTON
GUNSIGHT
MT DE BOER
Lakelse Lake
Skeena River
N
Williams Creek
0
10
kilometres

NAME OF STREAM WILLIAMS CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S.W. into N. end of Lakelse Lake.Rge. 5, Coast Dist.

POSITION 54 128 SW.

LENGTH 12.8 km WIDTH m DRAINAGE km^2

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m^2 SPAWNING AREA m^2

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- up to 6.4km from mouth
CHINOOK	- spawning grounds unknown
COHO	- up to 12.8km from mouth
CHUM	- near Highway #25 bridge.
PINK (ODD YEAR)	- spawning grounds unknown
PINK (EVEN YEAR)	" "
STEELHEAD	" "

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1949 - Scouring noted most years.

1951 - Stream bed damaged by flash floods. Advised that survey be made to assess feasibility of confining channel and controlling flow.

1952 - Stream bed deteriorated further.

1953 - Remedial stream work conducted to improve passage and flow conditions.

1954 - Numerous flash floods reported, 55.

1956 - Freshets caused stream bed damage 59,66,78

1958 - Freshet damage only partially surveyed.

continued.....

continued.

GENERAL REMARKS WILLIAMS CREEK

1961 - Some fish stripped for hatchery purposes.
1962 - Approx 10,000,000 eggs taken and about 4,100 fish allowed through the fence to spawn naturally.
1963 - Sockeye Creek escapements included in estimates this year.
1964 - Most sockeye released to spawn naturally.
1965 - B.C.Hydro clearing for power lines exposed trees alongside stream to wind effects and increased the possibility of their falling into stream.
1968 - Erosion noted 69-75
1974 - Considerable loss of sockeye spawn due to severe October floods.
1980 - Continued diversion of water into Blackwater Creek.

Log jams noted in lower reaches, but removed only when obstructing fish passage because they stabilize flow and reduce scouring.

Seasonal fluctuations in water levels--silty water.
Predators: bears and eagles.


× - Sockeye Spawning Grounds
▫ - Coho Spawning Grounds
- Marshy Area
- Road (Highway 25)
- Road Bridge
- Old Road Bridge
Williams
Creek
Lakelse
Lake
N
0
5.12
kilometres

Sketch of Williams Creek, 1971.

ESCAPEMENT RECORD FOR WILLIAMS CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947	7500		400			
48	7500		75		1500	
49	5200		300		N/O	
50	1400		400		N/O	
51	4000		250			
52	12000		1000		1000	
53	8500		300			
54	7500		400		N/O	
55	3500		400			
56	3500		750		N/O	
57	7500		1500			
58	7500		1500			
59	7500		N/I			
60	3500		N/O			
61	5500		N/O			
62	3500		400			
63	7500		N/O			
64	18000		400		75	
65	28761	25	400			
66	12731		1500			200
67	6586					
68	7500		1500			
69	3500		400			
70	3500		400			
71	1500		400			
72	1500		400	200		
73	1000		200			
74	1500		200			
75	1800		200			
76	1500		200			
77	1500		25			
78	1500		400			
79	9600		350			
80						
81	4000		300			
82						
83						
84						
85						

TIMING						
ARRIVE	L July		Oct	Aug		UNK
START	M Aug	Aug	Oct	E Sept	EAug-MSept	UNK
PEAK	LAug-MSept	Sept	MOct - MNov	M Sept	MAug-LSept	UNK
END	LSept-Oct	UNK	LNov - Dec	Oct	LAug-MOct	UNK

REMARKS N/I - not inspected


Wilson Creek
N
0
2
kilometres
NASS
MOUN
WM
WL
SKEENA RIVER
Wilson Creek
40
42
43
44
45
46
47
08
07
06
05
04
03
02
01
00
L 3952
L 2358
L 2357
L 2359
L 2618
L 842
L 2362
L 843
L 2363
L 2360
L 848
L 2351
BM 512
IR
BLK
5500
5000
4500
4000
3500
3000
2500
2000
1500

NAME OF STREAM WILSON CREEK

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows E. into Skeena River, N.E.of Cedarvale, Cassiar Dist.

POSITION 55 128 SE

LENGTH ____ km WIDTH ____ m DRAINAGE ____ km²

COMPOSITION: BEDROCK ____ BOULDER ____ COARSE ____ FINE ____

SILT & SAND ____ UNCLASSIFIED ____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA ____ m² SPAWNING AREA ____ m²

DISCHARGE (m³/s) ____

TEMPERATURE (°C) ____

BARRIERS OR POINTS OF DIFFICULT ASCENT 1977 - Narrow channel -- heavy rocks.
1978 - Log jams, narrow channel, heavy rocks at various locations.
1979 - Gravel bar at mouth. Stream drops 2-3m. recommend removal.
1980 - Impassable except in high water.
1981 - Recommend removal of gravel at mouth.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	- from mouth to approx 1km upstream
PINK (EVEN YEAR)	- " " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM ____

GENERAL REMARKS Physical conditions: Seasonal fluctuating water levels.
1974 - Heavy erosion silting and scouring on lower end of stream. Water levels normal.
1978 - 15-20% erosion and silting near mouth. Some scouring during Oct. flood.
1979 - Heavy erosion in lower half of stream due to fall flood. Heavy scouring-- large gravel deposit at mouth.
Suggest C.N.R. dig a deeper channel at confluence when they clear gravel build up under bridge next summer.



Sketch of Wilson Creek, 1971.

Wilson Creek
○ - Pink spawning grounds
□ - Coho spawning grounds
SKEENA RIVER
Cedarvale
N
0
.50
kilometres

ESCAPEMENT RECORD FOR WILSON CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68					200	
69			-		-	
70			75		400	
71			75		400	
72			75		400	
73			25		N/0	
74			-		N/0	
75			-		N/0	
76						
77						
78					200	
79					N/0	
80					N/0	
81					N/0	
82						
83						
84						
85						

TIMING						
ARRIVE			Oct		Aug	
START			M Oct		L Aug	
PEAK			Nov		Sept	
END			Dec		M Sept	

REMARKS


Zymagotitz River
0
10
kilometres
N
TREE FARM LICENCE 1
BLOCK 4
SKEENA MOUNTAINS
KITIMAT RANGES
FOREST
Kitsumkalum Lake
MT. ALLARD
1505
MT. GARLAND
2026
MAROON MTN
GLACIER PK.
1626
LEAN-TO MTN
1543
MT. VANARSDOLL
KITSELAS MTN
MT. HADDEN
MT. ZEUS
MT. NIELSON
MT. TRESTON
2041
MT. WILLIAM BROWN
MT. KENNEY
SLEEPING BEAUTY MTN
MT. MORRIS
MT. REMO
1954
KITSUMKALUM MTN
KITSUMKALUM MOUNTAIN RECREATION AREA
TERRACE
DIST. MUN.
WATERSHED RESERVE
COPPER MTN
Zymagotitz
Nelson
River
Exstew
Shames
Kalum
Maroon
Amsbury
Thornhill
750
1500

NAME OF STREAM ZYMAGOTITZ RIVER

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows S. into Skeena River, Rge.5., Coast Dist.

POSITION 54 128 SW.

LENGTH 29 km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Difficult though passable falls at 8.0km

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- lower 16.0km
COHO	- upper section and tributories.
CHUM	- lower 16.0km
PINK (ODD YEAR)	- " "
PINK (EVEN YEAR)	- " "
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1948 - Some erosion noted 68/70, 1970-78, 80,81

1960 - Silting in most years between 1960 and 81.

- Department of Northern Affairs installed an automatic flow recorder.

1961 - Logging in watershed 62,65.

1966 - Late coho and chums sighted

Steep rapids 19.3km from mouth. Swift and bouldery, difficult to inspect beyond 8.0km point.

Windfall and other debris monitored and removed as required. Subject to seasonal fluctuations in water levels.

Predators: bears.


Pine Lake
Pine Creek
Erlandson Creek
Molybenum Creek
Zymagotitz River
Skeena River
□ - Coho Spawning Grounds
▲ - Chum Spawning Grounds
○ - Pink Spawning Grounds
△ - Chinook Spawning Grounds
N - Passable Falls
N
0
5.12
kilometres

Sketch of Zymagotitz River, 1971.

ESCAPEMENT RECORD FOR ZYMAGOTITZ RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			750	750	750	
48			400	750	7500	
49			400	200	1500	
50			300	600	6000	
51			350	N/I	N/I	
52			300	100	7000	
53			N/I	200	3000	
54			N/I	200	3500	
55			N/I	75	3500	
56			750	75	750	
57			1500	200	7500	
58			1500	75	400	
59			N/I	N/I	N/I	
60			OBS		750	
61					3500	
62		25	25	25	400	
63			N/D	75	N/D	
64		25	50	200	N/O	
65			2500			
66			3500		200	
67			3500	75		
68		25	1500	N/O	750	
69		75	750	N/O	N/O	
70		75	400	N/O	N/O	
71		75	400		400	
72		75	400	400	1500	
73		200	400	400	750	
74		200	400	400	75	
75		200	400	200	75	
76		75	750	200	N/O	
77		75	400	200	25	
78		10	400	50	150	
79						
80		25	200	100	100	
81		20	400	N/O	50	
82						
83						
84						
85						

TIMING

ARRIVE		L July	Sept	Aug	Aug	
START		M Aug	M Oct	M Aug	MAug-ESept	
PEAK		L Aug	M Nov	L Aug	E-M Sept	
END		Sept	Dec	Sept	LSept - Oct	

REMARKS


N
Zymoetz River
0
20
kilometres
UNTAINS
Dorreen
Pacific
Usk
Lean-to Mtn
Terrace
Kitselas
Legate Creek
Kleanza Creek
Zymoetz
Williams Creek
Buildings
Lakelse Lake
River
Kitimat
Davies Ck
Mt Davies
Andesite Pk
Kitimat Ranges
Coast Mountains
Bulkley Ranges
Hazelton Mountains
Telkwa Range
Hudson Bay Mountain
McDonell
Kitseguecla R
Moricetown
Telkwa
Clore R
Burnie Lake
McBride
9039
5050
Mt Sir Robert
5850
6450
6150
6550
7450

NAME OF STREAM ZYMOETZ RIVER

CONSERVATION DISTRICT 8 STATISTICAL AREA 4

LOCATION OF MOUTH Flows N.W. into Skeena River. Rge. 5. Coast Dist.

POSITION

LENGTH km WIDTH m DRAINAGE km²

COMPOSITION: BEDROCK BOULDER COARSE FINE

SILT & SAND UNCLASSIFIED

PERCENT GRADIENT	
0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
>1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m^3/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Cascades at 6.4km and 9.6km.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	- in to McDonnell Lake and tributaries
CHINOOK	- throughout but concentrated in upper reaches.
COHO	- in small tributaries and secondary channels in upper section.
CHUM	- lower 8.0km
PINK (ODD YEAR)	- " "
PINK (EVEN YEAR)	- " "
STEELHEAD	- specific areas unknown

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1947 - Stream inspection incomplete.

1950 - Some fry were found emerging from gravel that had been covered by sand, but mortalities were high in areas of low water.

1951 - Flash floods considerably altered the lower 4.8km of the river.

1955 - Fish seen in riffle and jumping in lower river.

1957 - Columbia Cellulose Co. built access road alongside the river.

1958 - Crown Zellerbach logged in this area and agreements have been made to remove debris with the least impact and under Fisheries supervision.

continued.....

continued.

GENERAL REMARKS ZYMOETS RIVER

1960 - High water restricted coho sports fishing
     - Forestry Road provides increased access to stream for inspection purposes.
1961 - Poor sports fishery.
     - Some silting near mouth.
     - Sockeye inspection cancelled due to forest fire.
1961 - New highway bridge resulted in stabilization of lower river channel.
     - Columbia Cellulose road completed for 29km.
1964 - Good steelhead fishery 66,65.
     - Columbia Cellulose road opened to 48.2km and to date stream has been kept clear of debris.
1965 - Logging road in use for industrial and recreational purposes.
1966 - Severe flooding in October caused changes in stream pattern and flow. Road wash outs in Limonite Creek Region being repaired by Columbia Cellulose.
1967 - Rockslide 9.6km upstream prevented most coho and sockeye migrating upstream. Those surmounting slide were badly battered.
1968 - Channel blasted through rock slide 9.6km from mouth by Resource Development Branch in August.

silting and scouring common during high water 68-78
stream is very silty during most of the salmon runs and accurate estimates of escapements are impossible.
seasonal fluctuations in water levels.
predators: bears and eagles.

ESCAPEMENT RECORD FOR ZYMOETZ RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947	1500		heavy	average		
48		moderate	light	heavy	moderate	moderate
49						
50		average	poor	good	good	average
51			fair	OBS	very good	
52		light	good	light	very heavy	above ave.
53			light	average	good	heavy
54		average	average	N/O	good	above ave.
55		heavy	average		very good	average
56		average	average	light	light	average
57		average	above ave.	light	above ave.	light
58		average	above ave.	light	light	fair
59	average	average	average	average	average	
60			OBS			OBS
61	OBS		OBS			poor
62	350	25	75	25	N/O	light
63			average			average
64	250	25	1500	75	N/O	good
65	3000		400			good
66	350		1000			average
67	200		200			
68	3500	75	1500	N/O		good
69	1500	200	3500	N/O		good
70	1500	200	3500	N/O		average
71	1500	400	3500			average
72	1500	200	3500	400	35000	
73	2500	400	1500	400	1500	
74	2000	400	1500	400	400	
75	400	200	750	200	400	
76	400	200	1500	200	200	
77	1500	400	1500	200	200	
78	3500	200	3500	500	100	
79						
80	400	100	1000	25	1000	
81	100	600	700	N/O	50	
82						
83						
84						
85						

TIMING						
ARRIVE	July	Aug	L Aug	Aug	Aug	
START	M Aug	M Aug	Sept - EOct	E Sept	E Sept	
PEAK	LAug - ESept	L Aug	LOct - M Nov	M Sept	M Sept	Nov - Jan
END	Sept	Sept	Nov - Dec	LSept-Oct	L Sept-Oct	

REMARKS

APPENDIX 1


Burdick Creek
N
0
2
kilometres
FOREST
Hazelton Peak
Burdick
Creek
GM


Cedar River
Little Cedar River
Creek
River
WL
KITSUMKALUM
RIVER
SKEENA PROV FOREST
Dump
Private
85
84
83
82
81
80
79
78
77
76
07
08
09
10
11
12
13
14
15
L 1040
L 1039
L 1036
L 1037
L 1038
L 1035
L 442
L 441
L 439
L 440
L 438
L 437
L 436
L 435
L 4977
L 4976
L 4978
L 5125
L 5124
L 5122
L 1032
L 1034
L 1033
L 5120
L 5121
L 1044
L 4370
L 4368
L 5119
L 433
L 1043
L 4367
L 6737
L 4979
L 4980
1000
2000
1500
1800
1100
900
2500
500
600
640
N
0
2
kilometres


NASS
(HAZE
PRO
Clear Creek
River
KITSUMKALUM
RIVER
SKEENA PROV FOREST
Dump
Private
Rosswood
84
83
82
81
80
79
78
77
76
75
74
10
11
12
13
14
15
16
17
18
L 1040
L 1039
L 1036
L 1037
L 1038
L 1035
L 442
L 441
L 4977
L 5125
L 4976
L 439
L 4978
L 440
L 438
L 437
L 1032
L 5124
L 5122
L 5123
L 435
L 436
L 1034
L 1033
L 5120
L 1044
L 4370
L 4368
L 5121
L 433
L 5119
L 1043
L 4367
L 4979
L 432
1500
1100
2500
3000
2000
4500
1800
900
600
640
500
N
Clear Creek
0
2
kilometres


Clear Creek
N
0
2
kilometres
TFL RESERVE
Mount Couture
TREE FARM LICENCE 1 BLOCK 4
N A S S
R
WL
Clear
Cre
4500
2500
1500
1100
4000
3000
1500


Maroon Creek
Fiddler Creek
Fiddler Lake
Icefield
Hardscrabble Creek
Shannon Creek
Lourie Ck
ASS RANGES
ZELTON MOUNTAINS)
SKEENA
5597
5673
3500
4000
4500
5000
5300
6000
Fiddler Creek
0
2
kilometres


33
34
35
36
37
38
39
40
41
Hatchery Creek
3400
3900
4000
Gunsight Peak
Mount De Boer
TREE FARM
Schulbuckhand Creek
1900
2000
2200
Chis
N
Schulbuckhand Creek
0
2
kilometres

METRIC EQUIVALENTS

Length

centimeter (cm) = 0.394 in
meter (m) = 3.280 ft
meter (m) = 1.094 yd
kilometer (km) = 0.621 mi

inch (in) = 2.540 cm
foot (ft) = 0.305 m
yard (yd) = 0.914 m
mile (mi) = 1.609 km

Area

square centimeter (in^2) = 0.155 in^2
square meter (m^2) = 10.760 ft^2
square meter (m^2) = 1.196 yd^2
square kilometer (km^2) = 0.386 mi^2
hectare (ha) = 2.470 a

square inch (in^2) = 6.451 cm^2
square foot (ft^2) = 0.093 m^2
square yard (yd^2) = 0.836 m^2
square mile (mi^2) = 2.590 km^2
acre (a) = 0.405 ha

Volume

cubic centimeter (cm^3) = 0.061 in^3
liter (L) = 61.023 in^3
liter (L) = 0.035 ft^3
liter (L) = 0.264 U.S. gal
= 0.220 Imp. gal
cubic meter (m^3) = 35.315 ft^3
cubic meter (m^3) = 1.308 yd

cubic inch (in^3) = 16.387 cm^3
cubic inch (in^3) = 0.016 L
cubic foot (ft^3) = 0.028 m^3
cubic foot (ft^3) = 28.320 L
cubic yard (yd^3) = 0.765 m^3
U.S. gallon (gal) = 3.785 L
Imp. gallon (gal) = 4.546 L

Weight

gram (gm) = 0.035 oz
kilogram (kg) = 2.205 lb
kilogram (kg) = 0.001 ton (short)
tonne (t) = 1.103 ton (short)

ounce (oz) = 31.103 gm
pound (lb) = 0.373 kg
ton (short) = 907.180 kg
ton (short) = 0.907 t

Velocity

meter per second (m/s) = 3.280 ft/s
feet per second (ft/s) = 0.305 m/s

Discharge

cubic meter per second (m^3/s) = 35.315 ft^3/s
cubic foot per second (ft^3/s) = 0.028 m^3/s
cubic meter per second (m^3/s) = 15350.879 U.S. gal/min
= 13198.628 Imp. gal/min

Temperature

Degrees Centigrade (°C) = 5/9 (Degrees Fahrenheit - 32)
Degrees Fahrenheit (°F) = 9/5 (Degrees Centigrade) + 32